

**Opinion of the Scientific Panel on Food Additives,
Flavourings, Processing Aids and Materials in Contact with Food (AFC)
on a request from the Commission related to**

Butylbenzylphthalate (BBP) for use in food contact materials

Question N° EFSA-Q-2003-190

Adopted on 23 June 2005 by written procedure

SUMMARY

The Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC) has been asked to re-evaluate butylbenzylphthalate (BBP) for use in the manufacture of food contact materials.

Previously, a temporary Tolerable Daily Intake (t-TDI) of 0.1 mg/kg bw was set by the Scientific Committee for Food (SCF) based on the end-point of peroxisome proliferation in rodent liver. There is now a scientific consensus that liver peroxisome proliferation in rodents is not relevant for human risk assessment. The critical effects of BBP relate to reproduction. From the different studies available, the critical observations were as follows.

Decreases in epididymal spermatozoal concentrations have been reported at dose levels of 200 mg/kg bw/day and 2200 mg/kg bw/day, with a No Observed Adverse Effect Level (NOAEL) of 20 mg/kg bw/day and a Lowest Observed Adverse Effect Level (LOAEL) of 200 mg/kg bw/day. However, accompanying histopathological effects on the testes and adverse impact on fertility were only seen at 2200 mg/kg bw/day.

A NOAEL of 20 mg/kg bw/day and a LOAEL of 100 mg/kg bw/day were reported from a two-generation study, based on increased serum follicle stimulating hormone (FSH) concentrations in the F₀ parental males. Furthermore, in the same study all other examined end-points had a NOAEL of 100 mg/kg bw/day.

In another developmental toxicity study in the rat using a multiple dose design, foetotoxicity effects were observed. The overall benchmark dose, based on a 1% increase in abnormal testis location, was assessed at 95 mg/kg bw/day.

A multi-generation study including the F₂ generation had an overall NOAEL of 50 mg/kg bw/day, based on the reduction in anogenital distance in F₁ and F₂ offspring at 250 mg/kg bw/day.

Based on the current literature on BBP testicular toxicity, the Panel allocated a Tolerable Daily Intake (TDI) of 0.5 mg/kg bw, derived from a NOAEL of 50 mg/kg bw/day found in a multi-generation study and making use of an uncertainty factor of 100.

The limited available data on BBP concentration in foods and diets in UK and Denmark were used to provide an estimation of the dietary exposure. In the UK, mean and high (97.5 percentile) intakes of BBP from dietary sources were estimated to be respectively 0.008 and 0.020 mg/person/day in the adult population (equivalent to 0.1 and 0.3 µg/kg bw/day).

In a Danish study, estimated mean exposure ranged from 0.02 to 0.03 mg/day, i.e. 0.3 to 0.4 µg/kg bw/day considering a 70 kg adult. Based on the highest concentration of BBP determined, exposure at high percentiles was estimated as 0.32 mg/day equivalent to 4.5 µg/kg bw/day.

In another Danish study, the main dietary sources of exposure were estimated to be root crops (30%) and leaf crops (60%). The total daily oral intake at the regional level (Denmark) can be estimated to 1 µg/kg bw/day in the adults, 5.9 µg/kg bw/day in children aged 1 to 6 years and 2.4 µg/kg bw/day in children aged 7 to 14 years.

Based on the detection limit, intake from infant formulae was estimated 1.6 µg/kg bw/day in infants of less than 6 months and 0.7 µg/kg bw/day in infants of more than 6 months. For infants of more than 6 months, ready-to-use baby foods were also taken into account and the exposure was therefore estimated as less than 0.9 µg/kg bw/day.

The Panel noted that the dietary exposure to BBP (derived from packaging and other sources) may contribute up to about 1% of the TDI value.

KEY WORDS :

Butylbenzylphthalate, BBP, food contact materials, CAS n°85-68-7.

BACKGROUND

BBP may be present in food, either due to migration from food contact materials containing BBP or due to its widespread presence as an environmental contaminant which can be found in air, water, soil and food. BBP was evaluated by the Scientific Committee for Food (SCF) in 1988 and 1993 when a temporary Tolerable Daily Intake (t-TDI) for use in food contact materials was established based on the then most sensitive end-point of peroxisome proliferation in rodent liver (SCF, 1995). There is a scientific consensus that liver peroxisome proliferation in rodents is not a relevant endpoint for human risk assessment (IARC, 1995). The Panel has therefore been asked to re-evaluate BBP for use in food contact materials.

TERMS OF REFERENCE

The Commission asks EFSA to re-evaluate butylbenzylphthalate (BBP) for use in the manufacture of food contact materials.

ASSESSMENT

1. Chemistry

Identification of the substance

CAS No.:85-68-7

EINECS No.:201-622-7


IUPAC name:benzyl butyl phthalate

FCM Ref N°74560

Synonyms: 1,2-benzenedicarboxylic acid, butyl phenylmethyl ester; benzyl-n-butyl phthalate; phthalic acid, butyl benzyl ester.

Molecular formula: $C_{19}H_{20}O_4$

Structural formula:



Molecular weight:312.35

Purity/impurities

Degree of purity:>98.5 % (w/w)

Identity and percentage (w/w) of impurities:

< 1.0 % di-benzyl phthalate(CAS No. 523-31-9)

< 0.5 % benzyl benzoate(CAS No. 120-51-4)

< 0.5 % di-butyl phthalate(CAS No. 84-74-2)

<2 ppm α -chlorotoluene(CAS No. 100-44-7)

<2 ppm α - α -dichlorotoluene(CAS No. 98-87-3)

Physico-chemical properties

Physical state: oily liquid

Melting point: - 35 °C

Boiling point: 370 °C at 1010 hPa

Relative density: 1.120 g/cm³ at 20 °C

Vapour pressure:	$8.2 \pm 3.1 \times 10^{-5}$ hPa at 25 °C
Water solubility:	2.8 mg/l at 20 °C
Partition coefficient	n-octanol/water: $\log K_{ow}$ 4.84

2. Use

BBP is present in a large number of end products some of which are available for consumer use. The vast majority of BBP use, more than 90%, goes to plasticizing of PVC or other polymers. The BBP plasticized polymeric material has consumer and industrial uses such as flooring, sealants, and paints.

3. Exposure via food

A relatively small but significant use is in the food wrap or food packaging area. BBP has also been reported at low concentrations in baby equipment and children toys. However, in these products BBP probably occurs as by-product/impurity and has not been added intentionally (CSTEE, April, 1998).

No data on the levels of BBP in food in the EU attributable to migration from food contact materials have been submitted by the industry. Exposure assessment based on analytical determination of the total concentration of BBP in samples of foods or diets allow to assess overall dietary exposure, from packaging material and other sources.

An assessment of exposure to phthalates was performed based on the analysis of stored samples from the Total Diet Study conducted in the UK in 1993. Concentration data were combined with food consumption data from the National Diet and Nutrition Study of British Adults. Since phthalates are fat soluble, only those 10 food groups which make a major contribution to dietary fat intakes were selected for analysis. Among these, carcass meat, eggs, poultry and milk were analysed for individual concentrations of phthalates since these four food groups accounted for approximately 85% of the estimated dietary intake of total phthalates. Mean and high (97.5 percentile) intakes of BBP from these sources were estimated to be respectively 0.008 and 0.020 mg/person/day in the adult population equivalent to 0.1 and 0.3 $\mu\text{g}/\text{kg}$ bw/day for a 60 kg adult (MAFF, 1996).

In a Danish study (Petersen and Breindahl, 2000), BBP was analysed in 29 different meals collected by test persons during 24 hours. Results were normalised to a daily diet of 10 MJ of energy and to a body weight of 70 kg in order to estimate total daily exposure. Among the 29 samples, 8 were above the limit of determination. The estimated mean concentration of BBP in the diet varied according to the values assigned to the 21 samples which were under the limit of determination. Estimated mean exposure therefore ranged from 0.02 to 0.03 mg/day, i.e. 0.3 to 0.4 $\mu\text{g}/\text{kg}$ bw/day considering a 70 kg adult. Based on the highest concentration of BBP determined, exposure at high percentiles was estimated as 0.32 mg/day equivalent to 4.5 $\mu\text{g}/\text{kg}$ bw/day.

A further Danish assessment of BBP total dietary exposure based on estimated and measured concentrations in environmental compartments using the European Union System for the Evaluation of Substances (EUSES, a computer modeling program) was reported recently (Müller et al, 2003). The main dietary sources of exposure were estimated to be root crops (30%) and leaf crops (60%). These high contributions of vegetables to oral BBP exposure may indicate that the exposure estimates for the UK, as given above, might be underestimates of the actual exposure because the UK figures were only based on food from animal sources.

For Denmark, the total daily oral intake at the regional level was estimated to be 1 $\mu\text{g}/\text{kg}$ bw/day in the adults, 5.9 $\mu\text{g}/\text{kg}$ bw/day in children aged 1 to 6 years and 2.4 $\mu\text{g}/\text{kg}$ bw/day in children aged 7 to 14 years. These values do not differ in a large extent from the UK figures mentioned previously.

Exposure from infant formulae was estimated based on two scenarios: an infant of less than 6 months weighing 5.5 kg and ingesting 900 g/day of formulae and an infant of more than 6 months weighing 8 kg and ingesting 525 g/day of formulae. Based on the maximum measured concentration among 11 commercial products (0.01 mg/kg of wet weight), this would lead to an exposure of 1.6 µg/kg bw/day in the infant of less than 6 months and 0.7 µg/kg bw/day in the infant of more than 6 months. For infants of more than 6 months, exposure from ready-to-use baby foods was also taken into account, considering the daily consumption of one jar of 250 g containing 0.005 mg BBP/kg of wet weight (the maximum measured concentration among 11 commercial products). Total exposure from infant formulae and ready to use baby foods in infants aged more than 6 months was therefore estimated as 0.9 µg/kg bw/day.

In this study, total oral exposure was also assessed; it was estimated to be 0.97, 4.1, 5.9 and 2.4 µg/kg bw/day in adults, infants (6-12 months), children (1-6 years) and children (7-14 years), respectively. Total exposure from all sources including dermal and inhalation exposure was very similar: 1.0, 4.2, 6.0, and 2.5 µg/kg bw/day respectively, due to a very low estimated exposure from these sources via these two routes.

4. Toxicological evaluation

Introduction

The Panel did not carry out a new extensive risk assessment but took cognisance of the previous evaluations by the SCF and in particular considered the more recent BBP Risk Assessment Report (RAR), prepared for the European Union Existing Substances Regulation, 793/93, 2001 (Annex 1), and the comments of the Scientific Committee on Toxicity, Ecotoxicity and the Environment (CSTEE) on this RAR (Annex 2), in order to determine the most significant toxicological end-point for risk assessment. Based on this information, the Panel focused on the most sensitive toxicological end-points for the evaluation of BBP, taken from the reproduction/developmental toxicity studies with this substance.

The SCF expressed its opinion on BBP in 1988 (SCF, 1995) based on the following statement in the safety data sheet:

“Reviewing the evaluations on the phthalate esters and comparing the data with the data on di(2-ethylhexyl)phthalate, the phthalate ester studied most extensively, the Committee decided to use a safety factor of 1000 instead of 100 for calculating the TDI for butylbenzylphthalate. This means that a temporary TDI of 0.1 mg/kg bw was established and that a definite TDI can be established when an adequate teratogenicity study and a no-effect level for peroxisomal proliferation, based on liver weight, peroxisomal associated enzyme activities and ultra-microscopy of the liver are available.”

Studies considered by the AFC Panel

The following studies on reproduction and development toxicities considered by the Panel for the determination of a NOAEL which could be used as a basis for a TDI calculation are summarized below. Further details on these studies are given in Annex 1 (Risk Assessment Report) and Annex 2 (CSTEE opinion).

A 26-week oral study in F344 rats resulted in decreased testicular weights, atrophy of seminiferous tubules, a near total absence of mature sperm production and marked hypospermia in the epididymis at dose level of 2.5% in the diet (1417 mg/kg bw/day) (Hazelton, 1985).

In conjunction with the NTP carcinogenicity study on BBP (NTP, 1997) a 10-week modified mating study in rats was also performed. Decreases in epididymal spermatozoal concentrations have been reported at dose levels of 200 mg/kg bw/day and 2200 mg/kg bw/day, with a NOEL of 20 mg/kg bw/day and a LOAEL of 200 mg/kg bw/day. However, accompanying histopathological effects on the testes and adverse impact on fertility was only seen at 2200 mg/kg bw/day (NTP, 1997a).

A NOAEL of 20 mg/kg bw/day was reported from a two-generation study, based on increased serum follicle stimulating hormone (FSH) concentrations in the F₀ parental males (Nagao *et al.*, 2000). The corresponding LOAEL was 100 mg/kg bw/day. In the F₁ generation, the NOAEL was 100 mg/kg bw/day. Furthermore, in the same study all other examined end-points had a NOAEL of 100 mg/kg bw/day.

In another developmental toxicity study in the rat using a multiple dose design (Piersma *et al.*, 2000), BBP induced increased liver and kidney weights in dams, accompanied by liver enzyme increases in maternal serum. Observed foetotoxicity included increased resorptions, reduced foetal weights, increased incidence of skeletal anomalies, and reduced foetal testis weights in the presence of an increased incidence of retarded testicular descent. As embryotoxicity was found at lower dosages compared to observed maternal toxicity, BBP appeared to be a specifically embryotoxic compound. The overall benchmark dose was assessed at 95 mg/kg bw/day, based on a 1% increase in abnormal testis location.

In a multi-generation study (Tyl *et al.*, 2001, 2004), BBP was administered in the diet at 0, 750 (50), 3750 (250), and 11,250 mg/kg (750 mg/kg bw/day) ad libitum. Adult F₀ systemic toxicity and adult F₁ systemic and reproductive toxicity were present at 11,250 mg/kg (750 mg/kg bw/day). At 11,250 mg/kg, there were reduced F₁ and F₂ male anogenital distance (AGD) and body weights/litter during lactation, delayed acquisition of puberty in F₁ males and females, retention of nipples and areolae in F₁ and F₂ males, and male reproductive system malformations. At 3750 mg/kg (250 mg/kg bw/day), only reduced F₁ and F₂ offspring male AGD was present. There were no effects on parents or offspring at 750 mg/kg (50 mg/kg bw/day). The F₁ parental systemic and reproductive toxicity NOAEL was 3750 mg/kg. The offspring toxicity NOAEL was 750 mg/kg (50 mg/kg bw/day), based on the presence of reduced AGD in F₁ and F₂ males at birth at 3750 mg/kg, but no effects were observed on reproductive development, structures, or functions.

CONCLUSIONS

Based on all the available toxicological evidence, the Panel concludes that effects on reproduction and development are the most sensitive end-points on which to base the risk assessment for BBP. Previous reviews have identified as pivotal several rat reproduction studies conducted in the last decade, which gave NOAELs or LOAELs in the region of 20-100 mg/kg bw/day, with the critical effect being on male reproductive development.

Based on the current literature on BBP testicular toxicity and on the presence of reduced AGD in F1 and F2 males at birth at 250 mg/kg bw/day, (NOAEL 50 mg/kg bw/day) in the Tyl *et al.* study (2001, 2004), the Panel allocated a Tolerable Daily Intake (TDI) of 0.5 mg/kg bw, derived from a NOAEL of 50 mg/kg bw/day and making use of an uncertainty factor of 100.

The limited available data on BBP concentration in foods and diets in UK (1993) and Denmark (2003) were used to provide an estimation of the dietary exposure. In the UK, mean and high (97.5 percentile) intakes of BBP from dietary sources were estimated to be respectively 0.008 and 0.020 mg/person/day in the adult population (equivalent to 0.1 and 0.3 µg/kg bw/day).

In a Danish study (Petersen and Breindahl, 2000), BBP estimated mean exposure ranged from 0.02 to 0.03 mg/day, i.e. 0.3 to 0.4 µg/kg bw/day considering a 70kg adult. Based on the highest concentration of BBP determined, exposure at high percentiles was estimated as 0.32 mg/day equivalent to 4.5 µg/kg bw/day.

In another Danish study (Müller *et al.*, 2003), the main dietary sources of exposure were estimated to be root crops (30%) and leaf crops (60%). The total daily oral intake at the regional level (Denmark) can be estimated to 1 µg/kg bw/day in the adults, 5.9 µg/kg bw/day in children aged 1 to 6 years and 2.4 µg/kg bw/day in children aged 7 to 14 years.

Based on the detection limit, intake from infant formulae was estimated to be 1.6 µg/kg bw/day in infants of less than 6 months and 0.7 µg/kg bw/day in infants of more than 6 months. For infants of more than 6 months, ready-to-use baby foods were also taken into account and the exposure was therefore estimated as less than 0.9 µg/kg bw/day.

The Panel notes that the dietary exposure to BBP (derived from packaging and other sources) may contribute up to about 1% of the TDI value.

REFERENCES

- Agarwal DK, Maronpot JC, Lamb IV and Kluwe WM. (1985). Adverse effects of butyl benzyl phthalate on the reproductive and hematopoietic system of male rats. *Toxicology*, 35; 189-206.
- Albro PW and Thomas RO. (1974). *Biochem. Biophys. Acta*, 360; 380.
- Ashby J, Tinwell H, Lefevre PA, Odum J, Paton D, Millward SW, Tittensor S and Brooks AN. (1997). Normal sexual development of rats exposed to butyl benzyl phthalates from conception to weaning. *Regul. Toxicol. Pharmacol.*, 26; 102-118.
- BIBRA (1985). British Industrial Biological Research Association. Report No. 0495/1/84. A 21-days feeding study of butyl benzyl phthalate to rats: effects on the liver and liver lipids.
- BIBRA (1992). British Industrial Biological Research Association. Report No. 112/2/92. An investigation of the effect of n-butyl benzyl phthalate (BBP) on rat hepatic peroxisomes.
- Coldham NG, Dave M, Sivapathasundaram S, McDonnell DP, Connor C, Sauer MJ. (1997) Evaluation of recombinant yeast cell estrogen-screening assay. *Environ. Health Perspect.* 105 (7): 734-742.
- CSTEE (1998) Phthalate migration from soft PVC toys and child-care articles. EU Scientific Committee on Toxicity, Ecotoxicity and the Environment - Opinion expressed at the CSTEE third plenary meeting in Brussels, 24 April 1998.
- Hammond BG, Levinskas GL, Robinson EC and Johannsen FR. (1987). A review of the subchronic toxicity of butyl benzyl phthalate. *Toxicol. Ind. Health*, 3; 79-98.
- Hazelton Laboratories American Inc. (1985). Report submitted to the National Toxicology Program, Project Number 12307-02,3. Twenty-six week subchronic study and modified mating trial in F344 Rats Butyl Benzyl Phthalate- Final Report.
- IARC (1995). International Agency for Research on Cancer. Peroxisome Proliferation and its role in Carcinogenesis, views and expert opinions of an IARC Working Group Lyon, 7-11 Dec 1994, IARC Technical Report No.24 Lyon, 1995.
- Jobling S, Reynolds T, White R, Parker MG and Sumpter JP. (1995). A variety of environmentally persistent chemicals, including some phthalate plasticizers, are weakly estrogenic. *Environ. Health Perspect.*, 103; 582-587.
- Lake BG, Harris RA, Grasso P and Gangolli SD. (1978). Studies on the metabolism and biological effects of n-butyl benzyl phthalate in the rat, BIBRA, Report No. 232/78, Project No. BB-76-312. Monsanto study BB-76-032.
- MAFF (1996) Ministry of Agriculture, Fisheries and Food. Survey of plasticiser levels in food contact materials and in foods. *Food Surveillance Papers* No 21. 1996.
- Müller, A.M., Nielsen, A. and Ladefoged, O. (2003), Ministeriet for Fødevarer, Landbrug og Fiskeri Veterinær- og Fødevaredirektoratet. Human exposure to selected phthalates in Denmark, rapport 2003:15.
- Nagao Tetsuji, Ryo Ohta, Hideki Marumo, Tomoko Shindo, Shinsuke Yoshimura and Hiroshi Ono. (2000). Effect of butyl benzyl phthalate in Sprague-Dawley rats after gavage administration: a two-generation reproductive study. *Reprod Toxicology*, 14, 513-532.
- NTP. (1982). National Toxicology Program. NTP-80-25, NIH Publication No. 82-1769. Technical Report serie no. 213. Carcinogenesis bioassay of butyl benzyl phthalate in F344/N rats and B6C3F1 mice (feed study).

- NTP. (1989). National Toxicology Program. NTP Report No. 89-246. Final report Developmental toxicity evaluation of butyl benzyl phthalate administered in feed to CD rats on gestational day 6 to 15.
- NTP. (1990). National Toxicology Program. NTP report No. 90-114. Final report Developmental toxicity of butyl benzyl phthalate in CD-1-Swiss mice.
- NTP. (1991). National Toxicology Program. Report No. T-0035C. Final report on the reproductive toxicity of di-n-butyl phthalate in Sprague-Dawley rats.
- NTP. (1995). National Toxicology Program. Report No. 460, NIH publication No. 95-3376. Effects of dietary restriction on toxicology and carcinogenesis studies in F344/N rats and B6C3F1 mice.
- NTP. (1997). National Toxicology Program. Report No. 458, NIH publication No. 97-3374. Toxicology and carcinogenesis studies of butyl benzyl phthalate in F344/N rats (feed studies).
- Piersma AH, Verhoef A, te Biesebeek JD, Pieters MN and Slob W. (2000). Developmental toxicity of butyl benzyl phthalate in the rat using a multiple dose study design. *Reprod. Toxicology*, 14; 417-425.
- Petersen JH. and Breindhal T. (2000). Plasticizers in total diet samples, baby food and infant formulae. *Food Addit Contam*; 17 (2), 133-41.
- Risk Assessment Report on BBP according Regulation 739/93 (Existing Substances Regulation), NPCA, Norway, dated February 2002.
- SCF Report 33rd serie (1995). Opinion on butylbenzyl phthalate.
- Sharpe RM, Fisher JS, Millar MED MER, Jobling S and Sumpter JP. (1995). Gestational and lactational exposure of rats to xenoestrogens results in reduced testicular size and sperm production. *Environ. Health Perspect.*, 103; 1136-1143.
- Tyl RW, Myers CB and Marr MC. (2001). Two-generation reproductive toxicity evaluation of Butyl Benzyl Phthalate administered in the feed to CD (Sprague-Dawley) rats. RTI Project No. 65C-0726-200, RTI Protocol No. RTI-761.
- Tyl RW, Myers CB and Marr MC. (2004). Reproductive toxicity evaluation of Butyl Benzyl Phthalate in rats. *Reproductive Toxicology*, 18, 241-264.

SCIENTIFIC PANEL MEMBERS

Robert Anton, Sue Barlow, Dimitrios Boskou, Laurence Castle, Riccardo Crebelli, Wolfgang Dekant, Karl-Heinz Engel, Stephen Forsythe, Werner Grunow, Marina Heinonen, John Chr. Larsen, Catherine Leclercq, Wim Mennes, Maria Rosaria Milana, Iona Pratt, Ivonne Rietjens, Kjetil Svendsen, Paul Tobback, Fidel Toldrá.

ACKNOWLEDGEMENT

The Scientific Panel/Committee on Food Additives, Flavourings, Processing Aids and Materials in Contact with Food (AFC) wishes to thank Jean-Claude Lhuguenot for the preparation of this draft opinion.

ANNEX 1

Extract from the Risk Assessment Report on BBP Consolidated Final Report (dated February 2002)

Mutagenicity and carcinogenicity

BBP showed no evidence of mutagenicity in *Salmonell typhimurium* or mouse lymphoma cells. BBP did not induce sister chromatid exchanges (SCE) or chromosomal aberrations (CA) in CHO hamster cells. BBP induced morphological transformation in Syrian hamster embryo cells, but not in the BALB/3T3 cell transformation system. It did not induce sex-linked recessive lethals in *Drosophila melanogaster* or dominant lethal mutations in mice. Positive results were obtained in a mouse bone marrow test for SCE, however the responses were weak, and the SCE test was not repeated. For the induction of CA conflicting results were reported when different observations times were compared. No induction of micronucleus were reported in female rats after exposure to low doses of BBP (182.6 µg/kg bw/day during gestation and lactation). Based on the available data, and according to EU criteria, BBP should not be considered a mutagen.

Butyl benzyl phthalate was tested for carcinogenicity by oral administration in mice (NTP, 1982) and rats (NTP, 1982, 1997), including a dietary restriction study (NTP, 1997). No increase in the incidence of tumours were observed in mice. The results from the rat studies are summarised in Table 5. An increased incidence of mononuclear cell leukemias was reported in female rats at 12,000 ppm BBP. No significant increase was, however, found in two later studies with the same rat strain although a higher concentration was tested. An increased incidence of benign pancreatic tumours was seen in one conventional study in male rats, but not after dietary restriction. A marginally increased incidence of pancreatic adenomas occurred in female rats in a conventional study, but not after dietary restriction. Papillomas of the urinary bladder was marginally increased in female rats both in the conventionally study and after dietary restriction. Moreover, after dietary restriction and 32 months an increase in bladder carcinomas was found. The latter results are difficult to interpret as no historic controls are available. In one study in rats, butyl benzyl phthalate given prior to 7,12-dimethylbenz(a)anthracene inhibited mammary carcinogenesis. Thus, BBP appears to be a borderline case between no classification for carcinogenicity and. Cat. 3. However, due to the lack of genotoxic effects no classification is proposed.

Toxicity for reproduction and development

Reproductive effects of BBP and its major metabolites MBuP and MBeP in rats following oral administration both by gavage or in the diet have been investigated in studies of different duration (from 4 days to 26 weeks, and in 2-generation studies). The main effects reported include a decrease in the relative weight of testis, damage to the testis, epididymis, prostate, seminal vesicle and to reduced epididymal sperm concentrations, and at high BBP concentrations reduced fertility, in addition to increases in relative liver and kidney weights.

As regards the effects on fertility or reproductive organs following administration of BBP to rats in the diet (Tyl et al., 2001; Agarwal et al., 1985; NTP, 1997; Hammond et al., 1987) or via gavage (Piersma et al., 1995; Piersma et al., 2000; Lake et al., 1978; Nagao et al., 2000), reduced mating and

fertility indices, decreases in testis weight, histopathological changes in testis, and hormonal changes have been reported. These effects have in the majority of the studies been reported at BBP doses equal to (Hammond et al., 1987, 4-week diet study) or higher than those which have induced other effects, such as variations in absolute and relative weights of the liver and kidney and histopathological changes such as atrophy in the liver and pycnotic nuclei, acinar atrophy and slight fibrosis in the pancreas. Exceptions includes, when BBP is administered by gavage, a 14 days and a 4 days study in Sprague-Dawley rats (Lake et al., 1978), and a 28 days study in Cpb-WU rats (Piersma et al., 2000). In the Lake 14 days study, minimal testicular atrophy was reported in one of three animals examined at 480 mg/kg bw/day (Lake et al., 1978). In the 4 day study atrophic changes in the testis in 3 of 6 animals at 800 mg/kg bw/day of BBP were reported. In the 28 days study by Piersma et al., 2000 a decrease in testosterone level was reported from 450 mg/kg bw/day. Exceptions include, when BBP is administered in the diet, a 10 week fertility study (NTP, 1997). In this study a dose-related decrease in epididymal spermatozoa concentration compared to control animals was reported from 200 mg/kg bw/day (2800 ppm) ($p \leq 0.05$) of BBP and the NOAEL from this study was 20 mg/kg bw/day of BBP. When taking into account days of recovery in males in the 10 week NTP study (days from positive sperm plug to necropsy) in a covariate analysis of variance, on the epididymal spermatozoa concentration, the decrease in epididymal spermatozoa concentration at 2800 ppm was not statistically significant at the 5 % level compared to control animals ($p = 0.07$). However, the dose-dependent decrease in epididymal spermatozoa concentration was still evident. In the parallel 26-week oral toxicity study in rats where BBP was administered in the diet (NTP, 1997) the control value for epididymal spermatozoa concentration may not be valid due to a reported possible inadequate mincing of the cauda epididymis tissue from control animals. The NOAEL for reduced epididymal spermatozoa concentration and fertility in the 26 week study was 550 mg/kg bw/day. In a new 2-generation study (Tyl et al., 2001) significantly reduced mating and fertility indices were reported in F1 parents to make F2 offspring at 750 mg/kg bw/day. In the same study a significantly reduced relative and absolute paired ovaries and uterus weight was reported in F0 females. In adult F1 males a significant increase in reproductive tract malformations were reported (53.33 % compared to 0 % in controls). No increases in reproductive tract malformations were reported in females. Systemic toxicity reported at 750 mg/kg bw/day was limited to organ weight changes (liver, kidney) in males and females and histopathological lesions graded as minimal in females. The NOAEL for fertility was 250 mg/kg bw/day from this study. In another recent two-generation study (Nagao et al., 2000), increased serum FSH in F0 males was reported from 100 mg/kg bw/day, and at 500 mg/kg bw/day a decreased testosterone level. In F₁ males (18 weeks old) a decrease in testis, epididymis and ventral prostate weight, a decrease in testosterone and LH levels, and atrophy of the seminiferous tubules with decreased number of germ cells, and a decreased number of sperm in the epididymis were reported, accompanied with reduced body weight and an increased relative liver and kidney weight. No effect on fertility was reported in this study at any dose levels (20, 100 or 500 mg/kg bw/day). From the Nagao (2000) study, no NOAEL value could be derived for effects on fertility or the reproductive organs in parent animals. The NOAEL value for developmental effects was 20 mg/kg bw/day based on reduced body weight in male and female F1 offspring from 100 mg/kg bw/day. In the risk characterisation, a NOAEL value OF 20 mg/kg bw/day for effects on the reproductive organs could be used. This NOAEL value is based on a dose-dependent decrease in epididymal spermatozoa concentration from 200 mg/kg bw/day (NTP, 1997).

ANNEX 2

Extracts from CSTE opinion on the results of the Risk Assessment of BBP

Opinion expressed on 24-4-1998

Part on effects assessment

A 26-week oral study in F344 rats resulted in decreased testicular weights, atrophy of seminiferous tubules, a near total absence of mature sperm production and marked hypospermia in the epididymes at dose level of 2.5% in the diet (1417 mg/kg b/w./day). Relative liver weight was increased at the 0.83% level (470 mg/kg bw/day), but not at the 0.28% level (159 mg/kg bw/day) (Hazelton, 1985).

In an oral 90-day study in rats at dose levels of 151 to 1069 mg/kg bw/day, increased liver weights were seen at all dose levels in females and at the mid and high dose in males. The increased liver weights in females were only slight at the two lower doses. Cecal weights were also increased at all dose levels in females, but not in males. Histopathological lesions of the pancreas were observed in males at the mid and high doses. The liver of high dose males had small areas of necrosis. The LOAEL of increased liver and cecal weights in females was 171 mg/kg bw/day. The NOAEL for increased liver weight in males was 151 mg/kg bw/day, the NOAEL for pancreatic effects was 381 mg/kg bw/day (Hammond et al., 1987).

Biochemical evidence of liver peroxisomal proliferation in male F344 rats was determined at all dose levels in males and was apparent from a dietary level of 0.6% (approx. 300 mg/kg bw/day). Moderate electron microscopical evidence of liver peroxisomal proliferation was observed at the 2.5% dietary level (approx. 1250 mg/kg bw/day, this dose was the only one examined) (BIBRA, 1985). Also, a recent combined long-term and carcinogenicity assay identified a dose level of 300 mg/kg bw/day as the lowest effect level for peroxisome proliferation (NTP, 1997).

In a two-year bioassay with F344 rats, females at the high dose (1.2% dietary level, 1200 mg/kg bw/day) had increased incidence of mononuclear cell leukaemia (18/50 vs. 7/49 in control and also at the 0.6% dietary level). There were no increased incidence of tumours in males, but they had to be killed at week 29-30 due to internal haemorrhaging (NTP, 1982). No increased tumour incidences were observed in B6C3F1 mice fed for 103 weeks with 0, 0.6 (840 mg/kg bw/day) or 1.2% dietary levels (1680 mg/kg bw/day) of BBP (NTP, 1982). The increased incidence of mononuclear cell leukaemia in rats (NTP, 1982) was not seen in a new 2-year dietary study in F344/N rats (NTP, 1997). In this study, a statistically increased incidence of pancreatic neoplasms was found in males.

BBP is not genotoxic in a number of *in-vitro* tests or in a dominant lethal study in mice (Ashby et al., 1994).

BBP induces embryotoxicity and teratogenic effects at a maternally toxic dose of 750 mg/kg bw/day at days 7-9 or 13-15 of gestation, this was not due to decreased maternal food intake (Ema et al., 1991 and 1993). BBP is also embryotoxic and teratogenic at maternally toxic dietary levels of 1.25-2.0% (NTP, 1990). An oral developmental reproduction study with BBP in Wistar rats is ongoing at the TNO, The Netherlands (Protocol P 470839).

BBP has shown oestrogen-like effects *in-vitro* (Jobling et al., 1995). Sharpe et al. (1995) have reported that exposure of rats to BBP in drinking water at a level of 1000 µg/l (corresponding to a dose of 50 µg/kg bw/day) during pregnancy and lactation results in a reduction in testis size in the male offspring on postnatal day 90. However, administration of 1000 µg BBP/L to pregnant AP rats during gestation and lactation was not found to lead to changes in the sexual development of pups of either gender (Ashby et al., 1997). An as yet unpublished drinking water study performed by the TNO (sponsored by ECPI/CMA) with a similar design as the Sharpe et al.-Study, has also not been able to duplicate the Sharpe et al.-findings.

BBP given s.c. for three consecutive days up to an amount of 5 mg, did not induce increased uterine weight in prepubertal mice (Coldham et al., 1997).

The most sensitive clear critical effect reported in the repeated dose experiments was increased liver and cecal weights in female rats at the lowest dose of 171 mg/kg bw/day after 90-day feeding. No effects were found in males at 151 mg/kg bw/day (Hammond et al., 1987).

In a review document (BIBRA, 1992) it is stated that degeneration of testes were seen at 480 mg/kg bw/day/d, but the investigation itself was confidential. The study by Sharpe et al. (1995) have reported testicular toxic effects in the offspring at a very low dose (50 µg/kg bw/day/d), but Ashby et al. and ECPI/CMA (unpublished) failed to confirm these findings. It thus seems unwarranted to base a NOAEL on the findings of Sharpe et al. (1995).

Conclusion. The critical effect used for assignment of a NOAEL value for BBP are increased liver weight observed in an oral 90-day rat study (Hammond et al., 1987). This study did not identify a NOAEL value in female rats, the LOAEL value for the critical effect is 171 mg/kg bw/day. Because the LOAEL value is used and there was only a slight increase in liver weights, an additional uncertainty factor of 2 has to be incorporated.

Opinion expressed on 27-11-1998

Part on effects assessment

In conjunction with the NTP carcinogenicity study on BBP (NTP, 1997) a 10-week modified mating study in rats was also performed. At a dose of 200 mg BBP/kg bw/day (2800 ppm in feed) to a group of rats, the epididymal spermatozoal concentration was significantly less than the controls. The NOAEL for these effects was 20 mg/kg bw/day. For its reassessment the CSTE will apply this value instead of the LOAEL of 171 mg/kg bw/day (incorporating an additional uncertainty factor of 2), as was done in its opinion of 24 April 1998.