

**SVMA14-004**

**DOCUMENT M-CP, Section 3**

**DATA ON APPLICATION**

---

**Version history<sup>1</sup>**

<b>Date</b>	<b>Data points containing amendments or additions and brief description</b>	<b>Document identifier and version number</b>

<sup>1</sup> It is suggested that applicants adopt a similar approach to showing revisions and version history as outlined in SANCO/10180/2013 Chapter 4 How to revise an Assessment Report

---

---

## Table of Contents

<b>CP 3</b>	<b>DATA ON APPLICATION .....</b>	<b>4</b>
<b>CP 3.1</b>	<b>Field of Use Envisaged .....</b>	<b>4</b>
<b>CP 3.2</b>	<b>Effects on Harmful Organisms.....</b>	<b>4</b>
<b>CP 3.3</b>	<b>Details of Intended Use .....</b>	<b>5</b>
<b>CP 3.4</b>	<b>Application Rate and Concentration of Active Substance .....</b>	<b>7</b>
<b>CP 3.5</b>	<b>Method of Application .....</b>	<b>7</b>
<b>CP 3.6</b>	<b>Number and Timings of Applications and Duration of Protection.....</b>	<b>7</b>
<b>CP 3.7</b>	<b>Necessary Waiting Periods or Other Precautions to Avoid Phytotoxic Effects on Succeeding Crops .....</b>	<b>7</b>
<b>CP 3.8</b>	<b>Proposed Instructions for Use.....</b>	<b>7</b>

## **CP 3        DATA ON APPLICATION**

### **CP 3.1        Field of Use Envisaged**

SVMA14-004 is intended to be used in field as foliar spray in mixture with insecticides.

### **CP 3.2        Effects on Harmful Organisms**

SVMA14-004 contains Hydrolysed Proteins. Hydrolysed Proteins have no harmful effect over the insects and in particular over the fruit flies. The hydrolysed proteins are mainly nutritional products. They act as insect attractant in order to trap them or to make easier the contact with the insecticide.

### CP 3.3 Details of Intended Use

PPP (product name/code): SVMA14-004  
 Hydrolysed proteins: Hydrolysed proteins  
 Safener: -  
 Synergist: -  
 Applicant: PROALAN S.A.  
 Zone(s): Southern  
 Verified by MS: yes/no

Formulation type: Soluble concentrate [SL]  
 Conc. of as 1: 300 g/L  
 Conc. of safener: -  
 Conc. of synergist: -  
 Professional use: ☒  
 Non professional use: ☐

Field of use: Attractant

1	2	3	4	5	6	7	8	9	10	11	12	13	14
Use- No. <sup>(e)</sup>	Member state(s)	Crop and/ or situation  (crop destination / purpose of crop)	F, Fn, Fpn G, Gn, Gpn or I	Pests or Group of pests controlled  (additionally: developmental stages of the pest or pest group)	Application				Application rate			PHI (days)	Remarks:  e.g. g safener/synergist per ha <sup>(f)</sup>
					Method / Kind	Timing / Growth stage of crop & season	Max. number a) per use b) per crop/ season	Min. interval between applications (days)	L product / ha a) max. rate per appl. b) max. total rate per crop/season	kg as/ha a) max. rate per appl. b) max. total rate per crop/season	Water L/ha  min / max		
Zonal uses (field or outdoor uses, certain types of protected crops)													
1	Southern zone	Citrus	F	Mediterranean Fruit Fly <i>Ceratitis capitata</i>	Foliar application	#	#	#	a) 1.5  b) #	a) 0.45  b) #	#	#	To be used in mixture with insecticides # corresponds to the GAP of the insecticide used in mixture
2	Southern zone	Persimmon	F	Mediterranean Fruit Fly <i>Ceratitis capitata</i>	Foliar application	#	#	#	a) 1.5  b) #	a) 0.45  b) #	#	#	To be used in mixture with insecticides # corresponds to the GAP of the insecticide used in mixture

<b>Remarks table heading:</b>	(a)	e.g. wettable powder (WP), emulsifiable concentrate (EC), granule (GR)	(d)	Select relevant
	(b)	Catalogue of pesticide formulation types and international coding system CropLife International Technical Monograph n°2, 6th Edition Revised May 2008	(e)	Use number(s) in accordance with the list of all intended GAPs in Part B, Section 0 should be given in column 1
	(c)	g/kg or g/l	(f)	No authorization possible for uses where the line is highlighted in grey, Use should be crossed out when the notifier no longer supports this use.
<b>Remarks columns:</b>	1	Numeration necessary to allow references	7	Growth stage at first and last treatment (BBCH Monograph, Growth Stages of Plants, 1997, Blackwell, ISBN 3-8263-3152-4), including where relevant, information on season at time of application
	2	Use official codes/nomenclatures of EU Member States	8	The maximum number of application possible under practical conditions of use must be provided.
	3	For crops, the EU and Codex classifications (both) should be used; when relevant, the use situation should be described (e.g. fumigation of a structure)	9	Minimum interval (in days) between applications of the same product
	4	F: professional field use, Fn: non-professional field use, Fpn: professional and non-professional field use, G: professional greenhouse use, Gn: non-professional greenhouse use, Gpn: professional and non-professional greenhouse use, I: indoor application	10	For specific uses other specifications might be possible, e.g.: g/m³ in case of fumigation of empty rooms. See also EPPO-Guideline PP 1/239 Dose expression for plant protection products.
	5	Scientific names and EPPO-Codes of target pests/diseases/ weeds or, when relevant, the common names of the pest groups (e.g. biting and sucking insects, soil born insects, foliar fungi, weeds) and the developmental stages of the pests and pest groups at the moment of application must be named.	11	The dimension (g, kg) must be clearly specified. (Maximum) dose of a.s. per treatment (usually g, kg or L product / ha).
	6	Method, e.g. high volume spraying, low volume spraying, spreading, dusting, drench	12	If water volume range depends on application equipments (e.g. ULVA or LVA) it should be mentioned under "application: method/kind".
		Kind, e.g. overall, broadcast, aerial spraying, row, individual plant, between the plants	13	PHI - minimum pre-harvest interval
		- type of equipment used must be indicated.	14	Remarks may include: Extent of use/economic importance/restrictions

---

### **CP 3.4      Application Rate and Concentration of Active Substance**

SVMA14-004 is applied at a rate of 1.5 L product/ha corresponding to 450 g a.s./ha.

### **CP 3.5      Method of Application**

SVMA14-004 is applied by foliar spraying in mixture with insecticides. Therefore, the water volume varies depending on the insecticide used in mixture. Spray volume should be adjusted to suit the label instructions of the insecticide used in mixture. For more information please refer to point 3.3.

### **CP 3.6      Number and Timings of Applications and Duration of Protection**

SVMA14-004 is to be applied in mixture with insecticides. Therefore, the number and timing of application depend on the insecticide used in mixture. It should be adjusted to suit the label instructions of the insecticide used in mixture. For more information please refer to point 3.3.

### **CP 3.7      Necessary Waiting Periods or Other Precautions to Avoid Phytotoxic Effects on Succeeding Crops**

Not applicable. Hydrolysed Proteins is intended to be used on perennial crops only.

### **CP 3.8      Proposed Instructions for Use**

SVMA14-004 is to be applied in mixture with insecticides. Therefore, the instruction for use of the insecticide used in mixture should be followed strictly. Once the insecticide has been added into the spray tank, pour the required quantity of SVMA14-004, stir well and add the remaining amount of water. The mixture should be applied as soon as possible after mixing.