

# NUTREL

**Insect attract for the preparation of insecticide baits against olive fly, cherry fly, fruit fly (apple, pear, peach, apricot, plum, kiwi, berries), citrus fly, walnut fly - Mass trapping in trap bottles**

Registration Italian Ministry of Health n. 11502 - 22/11/2002

Composition	
PURE HYDROLYSED PROTEINS	g. 30 (=378 g/L)
Coformulants q.s. to	g. 100

CONTENTS: 0.250-0.5-1-5-10-20 liters

Batch n.

Authorization holder:

**SICIT 2000 SpA**

Via Arzignano, 80 36072

Chiampe (Vicenza)

Tel. 0444 450 946

Produced by:

**Althaller Italia S.r.l.**

Strada Com.le per Campagna, 5

20078 S. Colombano all'Ambro

Tel.00371-200257

**CAUTION:**

**HANDE WITH CARE**

Distributed by:

**DIACHEM SpA**

Via Tonale 15, 24061 Albano S.

Alessandro (BG). Tel 0363 355611

**CBC (Europe) S.r.l.**

Via E. Majorana, 2 - 20834 Nova

Milanese (MB) - Tel. 0362 - 365079

**PRECAUTIONARY STATEMENTS: General: keep away from children. Prevention: P220: Store away from food, feed, beverage. P270: Do not eat, drink or smoke when using this product. P280: Wear protective clothing and gloves.**

**SUPPLEMENTARY INFORMATION OH HAZARDS: EUH 401: To avoid risks to human health and environment, comply with the instructions for use.**

**SUPPLEMENTARY PROVISIONS: Keep the package well closed. Avoid any contact with skin, eyes and clothes. After handling and in case of contamination, accurately wash with water and soap.**

## MEDICAL INFORMATION

**In case of intoxication call a doctor for the usual first aid measures and show him the label of the product that has been mixed with NUTREL.**

## DIRECTIONS FOR USE

Treatments from the ground: use NUTREL at the concentration of 1.2%, in mixture with the insecticide at the rate usually recommended. Treat only a limited and well exposed to sun crown area, on alternate rows. Repeat the treatment in case of rain.

Preferably use air blasts at normal and high pressure to obtain a perfect atomization of the insecticide mixture. Spray 100-200 liters of water per hectare, depending on used equipment, orchard dimension and training system. Maximum rate: 2.4 L NUTREL/ha, equal to 907 g a.s./Ha.

Use insecticides with a persistence at least equal to the one of NUTREL (e.g.: formulations based on deltamethrin, lambda-cyhalothrin).

To ensure the success of the control, it is necessary to apply 2-3 treatments, with an interval of 10-30 days.

For a more precise application point out the appearance of the adults using adequate trap bottles.

Mass trapping: on average, place per hectare 90 traps with adequate lateral holes, containing 250 ml of NUTREL. Place the traps 1.5 m-high, on branches facing south, avoiding direct sunlight. Refill in case of evaporation of the attractive liquid. Maximum rate: 22.5 L NUTREL/ha, equal to 8.5 kg of a.s./ha.

## COMPATIBILITY

NUTREL has to be mixed with insecticides complying with the precautionary statements recommended for the product used in mixture.

**Warning:** in case of mixture with other formulations, respect the longer pre-harvest interval. Comply with the precautionary statements recommended for the most toxic products. In case of intoxication, inform the doctor on the mixture done.

Interrupt the treatment according to the directions for use recommended for the insecticides used in mixture.

## ATTENTION

Use exclusively for the uses and under the conditions reported in this label.

The user of the product is responsible for eventual damages deriving from its inappropriate use.

The compliance with all the directions reported in this label is a necessary condition to ensure the efficacy of the treatment and avoid damages to plant, people and animal.

**DO NOT APPLY WITH AERIAL EQUIPMENT.**

**DO NOT CONTAMINATE OTHER CROPS, FOOD AND BEVERAGE OR WATERCOURSES.**

**DO NOT OPERATE UPWIND.**

**DO NOT SELL THIS PRODUCT LOOSE.**

**THE DISPOSAL OF THE PACKAGE MUST COMPLY WITH THE LEGISLATION IN FORCE.**

**DO NOT RELEASE THE PACKAGE, COMPLETELY EMPTY, IN THE ENVIRONMENT.**

**DO NOT REUSE THE PACKAGE.**