

MALTA

The Report referred to in Article 9 of Directive 2003/99/EC

TRENDS AND SOURCES OF ZOONOSSES AND ZOOTIC AGENTS IN HUMANS, FOODSTUFFS, ANIMALS AND FEEDSTUFFS

including information on foodborne outbreaks,
antimicrobial resistance in zoonotic agents and some
pathogenic microbiological agents.

IN 2011

INFORMATION ON THE REPORTING AND MONITORING SYSTEM

Country: Malta

Reporting Year: 2011

| Laboratory name | Description | Contribution |
|---|-------------|--------------|
| National Veterinary Laboratory - Agriculture and Fisheries Regulation Division-Ministry for Rural Affairs and the Environment | | |

PREFACE

This report is submitted to the European Commission in accordance with Article 9 of Council Directive 2003/99/ EC*. The information has also been forwarded to the European Food Safety Authority (EFSA).

The report contains information on trends and sources of zoonoses and zoonotic agents in Malta during the year 2011 .

The information covers the occurrence of these diseases and agents in humans, animals, foodstuffs and in some cases also in feedingstuffs. In addition the report includes data on antimicrobial resistance in some zoonotic agents and commensal bacteria as well as information on epidemiological investigations of foodborne outbreaks. Complementary data on susceptible animal populations in the country is also given. The information given covers both zoonoses that are important for the public health in the whole European Community as well as zoonoses, which are relevant on the basis of the national epidemiological situation.

The report describes the monitoring systems in place and the prevention and control strategies applied in the country. For some zoonoses this monitoring is based on legal requirements laid down by the Community Legislation, while for the other zoonoses national approaches are applied.

The report presents the results of the examinations carried out in the reporting year. A national evaluation of the epidemiological situation, with special reference to trends and sources of zoonotic infections, is given. Whenever possible, the relevance of findings in foodstuffs and animals to zoonoses cases in humans is evaluated.

The information covered by this report is used in the annual Community Summary Report on zoonoses that is published each year by EFSA.

* Directive 2003/ 99/ EC of the European Parliament and of the Council of 12 December 2003 on the monitoring of zoonoses and zoonotic agents, amending Decision 90/ 424/ EEC and repealing Council Directive 92/ 117/ EEC, OJ L 325, 17.11.2003, p. 31

List of Contents

| | | |
|--------|--|----|
| 1 | ANIMAL POPULATIONS | 1 |
| 2 | INFORMATION ON SPECIFIC ZOOSE AND ZOONOTIC AGENTS | 10 |
| 2.1 | SALMONELLOSIS | 11 |
| 2.1.1 | General evaluation of the national situation | 11 |
| 2.1.2 | Salmonellosis in humans | 12 |
| 2.1.3 | Salmonella in foodstuffs | 19 |
| 2.1.4 | Salmonella in animals | 20 |
| 2.1.5 | Salmonella in feedingstuffs | 28 |
| 2.1.6 | Antimicrobial resistance in Salmonella isolates | 29 |
| 2.2 | CAMPYLOBACTERIOSIS | 35 |
| 2.2.1 | General evaluation of the national situation | 35 |
| 2.2.2 | Campylobacteriosis in humans | 35 |
| 2.2.3 | Antimicrobial resistance in Campylobacter isolates | 39 |
| 2.3 | LISTERIOSIS | 47 |
| 2.3.1 | General evaluation of the national situation | 47 |
| 2.3.2 | Listeriosis in humans | 48 |
| 2.4 | E. COLI INFECTIONS | 49 |
| 2.4.1 | General evaluation of the national situation | 49 |
| 2.5 | TUBERCULOSIS, MYCOBACTERIAL DISEASES | 49 |
| 2.5.1 | General evaluation of the national situation | 49 |
| 2.5.2 | Tuberculosis, mycobacterial diseases in humans | 50 |
| 2.5.3 | Mycobacterium in animals | 53 |
| 2.6 | BRUCELLOSIS | 55 |
| 2.6.1 | General evaluation of the national situation | 55 |
| 2.6.2 | Brucellosis in humans | 56 |
| 2.6.3 | Brucella in animals | 59 |
| 2.7 | YERSINIOSIS | 64 |
| 2.7.1 | General evaluation of the national situation | 64 |
| 2.7.2 | Yersiniosis in humans | 65 |
| 2.8 | TRICHINELLOSIS | 66 |
| 2.8.1 | General evaluation of the national situation | 66 |
| 2.8.2 | Trichinellosis in humans | 67 |
| 2.8.3 | Trichinella in animals | 70 |
| 2.9 | ECHINOCOCCOSIS | 72 |
| 2.9.1 | General evaluation of the national situation | 72 |
| 2.9.2 | Echinococcosis in humans | 73 |
| 2.10 | TOXOPLASMOSIS | 76 |
| 2.10.1 | General evaluation of the national situation | 76 |
| 2.10.2 | Toxoplasmosis in humans | 76 |
| 2.10.3 | Toxoplasma in animals | 79 |

| | | |
|--------|---|-----|
| 2.11 | RABIES | 80 |
| 2.11.1 | General evaluation of the national situation | 80 |
| 2.11.2 | Rabies in humans | 81 |
| 2.12 | STAPHYLOCOCCUS INFECTION | 82 |
| 2.12.1 | General evaluation of the national situation | 82 |
| 2.13 | Q-FEVER | 82 |
| 2.13.1 | General evaluation of the national situation | 82 |
| 3 | INFORMATION ON SPECIFIC INDICATORS OF ANTIMICROBIAL | 83 |
| 3.1 | ESCHERICHIA COLI, NON-PATHOGENIC | 84 |
| 3.1.1 | General evaluation of the national situation | 84 |
| 3.1.2 | Antimicrobial resistance in Escherichia coli, non-pathogenic | 85 |
| 3.2 | ENTEROCOCCUS, NON-PATHOGENIC | 89 |
| 3.2.1 | General evaluation of the national situation | 89 |
| 3.2.2 | Antimicrobial resistance in Enterococcus, non-pathogenic isolates | 89 |
| 4 | INFORMATION ON SPECIFIC MICROBIOLOGICAL AGENTS | 96 |
| 4.1 | ENTEROBACTER SAKAZAKII | 97 |
| 4.1.1 | General evaluation of the national situation | 97 |
| 4.1.2 | Cronobacter in foodstuffs | 98 |
| 4.2 | HISTAMINE | 99 |
| 4.2.1 | General evaluation of the national situation | 99 |
| 4.3 | STAPHYLOCOCCAL ENTEROTOXINS | 100 |
| 4.3.1 | General evaluation of the national situation | 100 |
| 5 | FOODBORNE OUTBREAKS | 101 |

1. ANIMAL POPULATIONS

The relevance of the findings on zoonoses and zoonotic agents has to be related to the size and nature of the animal population in the country.

A. Information on susceptible animal population

Sources of information

National Livestock Database, National Veterinary Laboratory of the Agriculture and fisheries Regulation Division (CA)

Dates the figures relate to and the content of the figures

2011

Geographical distribution and size distribution of the herds, flocks and holdings

The data covers both the islands of Malta and Gozo. Malta's location is set in the heart of the Mediterranean Sea, half way between Sicily and North Africa, the Maltese archipelago consists of five islands: Malta, Gozo, Comino and two uninhabited islands, Cominotto and Filfla. The size of Malta is about 27 km long by 14.5 km width. Most of the farms for all species are found mainly on the main island i.e. Malta.

Table Susceptible animal populations

* Only if different than current reporting year

| Animal species | Category of animals | Number of herds or flocks | | Number of slaughtered animals | | Livestock numbers (live animals) | | Number of holdings | |
|-------------------------|---|---------------------------|-------|-------------------------------|-------|----------------------------------|-------|--------------------|-------|
| | | Data | Year* | Data | Year* | Data | Year* | Data | Year* |
| Cattle (bovine animals) | meat production animals | 326 | | 4738 | | 310 | | 326 | |
| | dairy cows and heifers | 125 | | 8953 | | 1638 | | 125 | |
| | calves (under 1 year) | 326 | | 4692 | | 2322 | | 326 | |
| | mixed herds | 125 | | 13564 | | 4270 | | 125 | |
| | - in total | 326 | | 14126 | | 4270 | | 326 | |
| Deer | farmed - in total ¹⁾ | | | | | | | | |
| Ducks | meat production flocks ²⁾ | | | | | | | | |
| | parent breeding flocks ³⁾ | | | | | | | | |
| | grandparent breeding flocks ⁴⁾ | | | | | | | | |
| | elite breeding flocks ⁵⁾ | | | | | | | | |
| | breeding flocks, unspecified - in total ⁶⁾ | | | | | | | | |
| | laying ducks ⁷⁾ | | | | | | | | |
| | mixed flocks/holdings ⁸⁾ | | | | | | | | |

Table Susceptible animal populations

| Animal species | Category of animals | Number of herds or flocks | | Number of slaughtered animals | | Livestock numbers (live animals) | | Number of holdings | |
|----------------------|---|---------------------------|-------|-------------------------------|-------|----------------------------------|-------|--------------------|-------|
| | | Data | Year* | Data | Year* | Data | Year* | Data | Year* |
| Ducks | - in total ⁹⁾ | | | | | | | | |
| Gallus gallus (fowl) | breeding flocks for egg production line - in total ¹⁰⁾ | | | | | | | | |
| | breeding flocks for meat production line - in total ¹¹⁾ | | | | | | | | |
| | breeding flocks, unspecified - in total ¹²⁾ | | | | | | | | |
| | elite breeding flocks for egg production line ¹³⁾ | | | | | | | | |
| | elite breeding flocks for meat production line ¹⁴⁾ | | | | | | | | |
| | elite breeding flocks, unspecified - in total ¹⁵⁾ | | | | | | | | |
| | parent breeding flocks for egg production line ¹⁶⁾ | | | | | | | | |
| | parent breeding flocks for meat production line ¹⁷⁾ | | | | | | | | |
| | parent breeding flocks, unspecified - in total ¹⁸⁾ | | | | | | | | |
| | grandparent breeding flocks for egg production line ¹⁹⁾ | | | | | | | | |
| | grandparent breeding flocks for meat production line ²⁰⁾ | | | | | | | | |
| | grandparent breeding flocks, unspecified - in total ²¹⁾ | | | | | | | | |
| | laying hens | 102 | | | | 255292 | | 38 | |

Table Susceptible animal populations

| Animal species | Category of animals | Number of herds or flocks | | Number of slaughtered animals | | Livestock numbers (live animals) | | Number of holdings | |
|----------------------|--|---------------------------|-------|-------------------------------|-------|----------------------------------|-------|--------------------|-------|
| | | Data | Year* | Data | Year* | Data | Year* | Data | Year* |
| Gallus gallus (fowl) | broilers | 567 | | | | 2385716 | | 74 | |
| Geese | meat production flocks ²²⁾ | | | | | | | | |
| | parent breeding flocks ²³⁾ | | | | | | | | |
| | grandparent breeding flocks ²⁴⁾ | | | | | | | | |
| | elite breeding flocks ²⁵⁾ | | | | | | | | |
| | breeding flocks, unspecified - in total ²⁶⁾ | | | | | | | | |
| | laying geese ²⁷⁾ | | | | | | | | |
| | mixed flocks/holdings ²⁸⁾ | | | | | | | | |
| Goats | meat production animals | 0 | | 0 | | 0 | | 0 | |
| | animals under 1 year | 294 | | 563 | | 937 | | 294 | |
| | animals over 1 year | 778 | | 286 | | 4023 | | 778 | |
| | milk goats | 778 | | 233 | | 3662 | | 778 | |
| | mixed herds | 861 | | 849 | | 4960 | | 861 | |
| | - in total | 861 | | 849 | | 4960 | | 861 | |

Table Susceptible animal populations

| Animal species | Category of animals | Number of herds or flocks | | Number of slaughtered animals | | Livestock numbers (live animals) | | Number of holdings | |
|----------------|---|---------------------------|-------|-------------------------------|-------|----------------------------------|-------|--------------------|-------|
| | | Data | Year* | Data | Year* | Data | Year* | Data | Year* |
| Pigs | breeding animals | 0 | | 4121 | | | | 0 | |
| | fattening pigs | 19 | | 34470 | | | | 19 | |
| | breeding animals - unspecified - sows and gilts | 0 | | 0 | | | | 0 | |
| | mixed herds | 138 | | 0 | | | | 138 | |
| | - in total | 157 | | 38591 | | | | 157 | |
| Reindeers | farmed - in total ²⁹⁾ | | | | | | | | |
| Sheep | meat production animals | 0 | | 0 | | 0 | | | |
| | animals under 1 year (lambs) | 598 | | 1345 | | 2623 | | | |
| | animals over 1 year | 1327 | | 1733 | | 9401 | | | |
| | milk ewes | 1327 | | 1633 | | 9022 | | | |
| | mixed herds | 1327 | | 3178 | | 12024 | | | |
| | - in total | 1403 | | 3178 | | 12024 | | | |
| Turkeys | meat production flocks ³⁰⁾ | | | | | | | | |
| | parent breeding flocks ³¹⁾ | | | | | | | | |

Table Susceptible animal populations

| Animal species | Category of animals | Number of herds or flocks | | Number of slaughtered animals | | Livestock numbers (live animals) | | Number of holdings | |
|----------------|--|---------------------------|-------|-------------------------------|-------|----------------------------------|-------|--------------------|-------|
| | | Data | Year* | Data | Year* | Data | Year* | Data | Year* |
| Turkeys | grandparent breeding flocks ³²⁾ | | | | | | | | |
| | elite breeding flocks ³³⁾ | | | | | | | | |
| | breeding flocks, unspecified - in total ³⁴⁾ | | | | | | | | |
| Wild boars | farmed - in total ³⁵⁾ | | | | | | | | |

Comments:

- ¹⁾ no deer are farmed in the Maltese Islands
- ²⁾ no ducks are farmed in the Maltese Islands
- ³⁾ no ducks are farmed in the Maltese Islands
- ⁴⁾ no ducks are farmed in the Maltese Islands
- ⁵⁾ no ducks are farmed in the Maltese Islands
- ⁶⁾ no ducks are farmed in the Maltese Islands
- ⁷⁾ no ducks are farmed in the Maltese Islands
- ⁸⁾ no ducks are farmed in the Maltese Islands
- ⁹⁾ no ducks are farmed in the Maltese Islands
- ¹⁰⁾ No breeding flocks are farmed in the Maltese Islands
- ¹¹⁾ flocks are farmed in the Maltese Islands
- ¹²⁾ flocks are farmed in the Maltese Islands
- ¹³⁾ flocks are farmed in the Maltese Islands

Table Susceptible animal populations

Comments:

- ¹⁴⁾ flocks are farmed in the Maltese Islands
- ¹⁵⁾ flocks are farmed in the Maltese Islands
- ¹⁶⁾ flocks are farmed in the Maltese Islands
- ¹⁷⁾ flocks are farmed in the Maltese Islands
- ¹⁸⁾ flocks are farmed in the Maltese Islands
- ¹⁹⁾ flocks are farmed in the Maltese Islands
- ²⁰⁾ flocks are farmed in the Maltese Islands
- ²¹⁾ flocks are farmed in the Maltese Islands
- ²²⁾ No geese are farmed on the Maltese Islands
- ²³⁾ No geese are farmed on the Maltese Islands
- ²⁴⁾ No geese are farmed on the Maltese Islands
- ²⁵⁾ No geese are farmed on the Maltese Islands
- ²⁶⁾ No geese are farmed on the Maltese Islands
- ²⁷⁾ No geese are farmed on the Maltese Islands
- ²⁸⁾ No geese are farmed on the Maltese Islands
- ²⁹⁾ No Reindeers are farmed in the Maltese Islands
- ³⁰⁾ No Turkeys are farmed in the Maltese Islands
- ³¹⁾ No Turkeys are farmed in the Maltese Islands
- ³²⁾ No Turkeys are farmed in the Maltese Islands
- ³³⁾ No Turkeys are farmed in the Maltese Islands
- ³⁴⁾ No Turkeys are farmed in the Maltese Islands
- ³⁵⁾ No Wild boar occurs in the Maltese Islands

Table Susceptible animal populations

2. INFORMATION ON SPECIFIC ZOOONOSES AND ZOONOTIC AGENTS

Zoonoses are diseases or infections, which are naturally transmissible directly or indirectly between animals and humans. Foodstuffs serve often as vehicles of zoonotic infections. Zoonotic agents cover viruses, bacteria, fungi, parasites or other biological entities that are likely to cause zoonoses.

2.1 SALMONELLOSIS

2.1.1 General evaluation of the national situation

A. General evaluation

History of the disease and/or infection in the country

Salmonella has been on the rise in Malta again since 2008. This follows a trough from 1999 - 2007. Majority of cases are S. Enteritidis and Typhimurium.

National evaluation of the recent situation, the trends and sources of infection

Most cases of Salmonella originate from households, including outbreaks (30-45 %). the sources are usually eggs and chicken meat. Information about salmonella in pork is not available.

Recent actions taken to control the zoonoses

Control of layers and broilers with culling for positive flocks of S. Enteritidis and S. Typhimurium. Heat treatment of eggs from infected layer flocks is also carried out.

2.1.2 Salmonellosis in humans

A. Salmonellosis in humans

Reporting system in place for the human cases

Physician and laboratory surveillance in place.

Case definition

Clinical picture with isolation of *Salmonella* from clinical specimens.

A probable case has an epidemiological link with a confirmed case.

Diagnostic/analytical methods used

Culture methods and serology.

Notification system in place

Statutorily notifiable.

History of the disease and/or infection in the country

Salmonella has been on the rise in Malta again since 2008. This follows a trough from 1999 - 2007.

National evaluation of the recent situation, the trends and sources of infection

Most cases of *Salmonella* originate from households, including outbreaks (30-45 %). The sources are usually eggs and chicken meat. Information about *salmonella* in pork is not available.

Table Salmonella in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. | Autochtho n cases | Autochtho n Inc. | Imported cases | Imported Inc. | Unknown status |
|-------------------------------|-------|------------|----------------------|---------------------|-------------------|------------------|-------------------|
| Salmonella | 74 | 17.75 | 74 | 17.75 | 0 | 0 | 0 |
| S. Enteritidis | 50 | 12 | 50 | 12 | 0 | 0 | 0 |
| S. Typhimurium | 24 | 5.75 | 24 | 5.75 | 0 | 0 | 0 |

Table Salmonella in humans - Age distribution

| Age distribution | S. Enteritidis | | | S. Typhimurium | | | Salmonella spp. | | | S. Bochum | | | S. Concord | | |
|--------------------|----------------|----|----|----------------|---|----|-----------------|----|---|-----------|---|---|------------|---|---|
| | All | M | F | All | M | F | All | M | F | All | M | F | All | M | F |
| <1 year | 3 | 2 | 1 | 1 | 0 | 1 | 1 | 1 | 0 | 1 | 0 | 1 | 2 | 1 | 1 |
| 1 to 4 years | 11 | 4 | 7 | 8 | 5 | 3 | 6 | 3 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5 to 14 years | 13 | 9 | 4 | 4 | 1 | 3 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 4 | 1 | 3 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 to 44 years | 6 | 3 | 3 | 2 | 0 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 45 to 64 years | 5 | 3 | 2 | 1 | 1 | 0 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 65 years and older | 7 | 0 | 7 | 6 | 2 | 4 | 3 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| Age unknown | 1 | 0 | 1 | 2 | 0 | 2 | 5 | 3 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : | 50 | 22 | 28 | 24 | 9 | 15 | 20 | 11 | 9 | 1 | 0 | 1 | 2 | 1 | 1 |

| Age distribution | S. Elomrane | | | S. Hadar | | | S. Haifa | | | S. Infantis | | | S. Kedougou | | |
|------------------|-------------|---|---|----------|---|---|----------|---|---|-------------|---|---|-------------|---|---|
| | All | M | F | All | M | F | All | M | F | All | M | F | All | M | F |
| <1 year | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 1 | 1 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 to 44 years | 1 | 0 | 1 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 |
| 45 to 64 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 0 | 0 | 0 | 0 |

Table Salmonella in humans - Age distribution

| Age distribution | S. Elomrane | | | S. Hadar | | | S. Haifa | | | S. Infantis | | | S. Kedougou | | |
|--------------------|-------------|---|---|----------|---|---|----------|---|---|-------------|---|---|-------------|---|---|
| | All | M | F | All | M | F | All | M | F | All | M | F | All | M | F |
| 65 years and older | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 1 |
| Age unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : | 1 | 0 | 1 | 1 | 0 | 1 | 4 | 2 | 2 | 5 | 4 | 1 | 1 | 0 | 1 |

| Age distribution | S. Kentucky | | | S. Livingstone | | | S. London | | | S. Mkamba | | | S. Newport | | |
|--------------------|-------------|---|---|----------------|---|---|-----------|---|---|-----------|---|---|------------|---|---|
| | All | M | F | All | M | F | All | M | F | All | M | F | All | M | F |
| <1 year | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 to 44 years | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 45 to 64 years | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 65 years and older | 2 | 2 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Age unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 2 | 0 | 2 |
| Total : | 4 | 3 | 1 | 2 | 0 | 2 | 1 | 1 | 0 | 1 | 0 | 1 | 2 | 0 | 2 |

Table Salmonella in humans - Age distribution

| Age distribution | S. Typhi | | | S. Virchow | | | S. enterica subsp. enterica | | | S. enterica subsp. salamae | | | Salmonella spp., unspecified | | |
|--------------------|----------|---|---|------------|---|---|-----------------------------|---|---|----------------------------|---|---|------------------------------|---|---|
| | All | M | F | All | M | F | All | M | F | All | M | F | All | M | F |
| <1 year | 0 | 0 | 0 | 3 | 2 | 1 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 2 |
| 5 to 14 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 to 44 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 45 to 64 years | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| 65 years and older | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 |
| Age unknown | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : | 2 | 1 | 1 | 4 | 2 | 2 | 3 | 1 | 2 | 1 | 1 | 0 | 4 | 0 | 4 |

Table Salmonella in humans - Seasonal distribution

| Seasonal Distribution | S. Enteritidis | S. Typhimurium | Salmonella spp. | S. Bochum | S. Concord | S. Elomrane | S. Hadar | S. Haifa | S. Infantis | S. Kedougou | S. Kentucky | S. Livingstone | S. London | S. Mkamba | S. Newport |
|-----------------------|----------------|----------------|-----------------|-----------|------------|-------------|----------|----------|-------------|-------------|-------------|----------------|-----------|-----------|------------|
| Months | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases | Cases |
| January | 4 | 1 | 0 | | | | | | | | | 1 | | | |
| February | 0 | 1 | 0 | | | | | | | | | | | | |
| March | 2 | 1 | 1 | | 2 | | | | | | 3 | | | | |
| April | 5 | 1 | 5 | | | | | | 1 | | | | | | |
| May | 2 | 2 | 1 | | | | | | | | | | | | |
| June | 7 | 2 | 0 | | | | | | | | | | | | |
| July | 4 | 2 | 2 | | | | | | | | | | | | |
| August | 7 | 1 | 2 | | | 1 | | 1 | 2 | 1 | | 1 | | | |
| September | 5 | 7 | 6 | | | | 1 | 0 | 2 | | | | | | 2 |
| October | 8 | 2 | 1 | 1 | | | | 1 | | | 1 | | 1 | 1 | |
| November | 1 | 1 | 0 | | | | | 2 | | | | | | | |
| December | 5 | 3 | 2 | | | | | | | | | | | | |
| not known | 0 | 0 | 0 | | | | | | | | | | | | |
| Total : | 50 | 24 | 20 | 1 | 2 | 1 | 1 | 4 | 5 | 1 | 4 | 2 | 1 | 1 | 2 |

| Seasonal Distribution | S. Typhi | S. Virchow | S. enterica subsp. enterica | S. enterica subsp. salamae | Salmonella spp., unspecified |
|-----------------------|----------|------------|-----------------------------|----------------------------|------------------------------|
| Months | Cases | Cases | Cases | Cases | Cases |
| January | | | | | |

Table Salmonella in humans - Seasonal distribution

| Seasonal Distribution Months | S. Typhi | S. Virchow | S. enterica subsp. enterica | S. enterica subsp. salamae | Salmonella spp., unspecified |
|---------------------------------|----------|------------|-----------------------------------|----------------------------------|---------------------------------|
| | Cases | Cases | Cases | Cases | Cases |
| February | 1 | | | | |
| March | | | | | |
| April | | | | | 1 |
| May | | | | | |
| June | | 1 | | | |
| July | | 2 | | | |
| August | | | | 1 | |
| September | | 1 | | | |
| October | | | 3 | | |
| November | 1 | | | | 3 |
| December | | | | | |
| not known | | | | | |
| Total : | 2 | 4 | 3 | 1 | 4 |

2.1.3 Salmonella in foodstuffs

A. Salmonella spp. in eggs and egg products

Monitoring system

Sampling strategy

there are no egg packing center each farmer packs his own eggs. Eggs were analysed on two occasions as re-confirmatory according to the Commission Regulation 1237/2007.

Diagnostic/analytical methods used

Raw material for egg products (at production plant)

Bacteriological method: ISO 6579:2002

2.1.4 Salmonella in animals

A. Salmonella spp. in Gallus Gallus - breeding flocks

Monitoring system

Sampling strategy

Breeding flocks (separate elite, grand parent and parent flocks when necessary)

There are no breeding flocks in Malta

B. Salmonella spp. in Gallus Gallus - broiler flocks

Monitoring system

Sampling strategy

Broiler flocks

All broiler flocks are sampled irrespective of their capacity. currently the sampling and analysis of both official controls and on behalf of the business operator are being carried out by the Competent Authority; namely the National Veterinary Laboratory

Frequency of the sampling

Broiler flocks: Before slaughter at farm

__2-3__ weeks prior to slaughter

Type of specimen taken

Broiler flocks: Before slaughter at farm

Socks/ boot swabs

Methods of sampling (description of sampling techniques)

Broiler flocks: Before slaughter at farm

The number of boot swabs taken per holding depends on the capacity of the farm. Six pairs of boot swabs are taken if the farms has a capacity of over 50,000. Four boot swabs are taken when there is a capacity between 10,000 - 50,000. Two pairs of boot swabs are taken when the capacity is less than 10,00. Two pairs of boot swabs are taken from each house sampled.

Case definition

Broiler flocks: Before slaughter at farm

A positive case is when Salmonella enteritidis and Salmonella typhimurium are isolated.

Diagnostic/analytical methods used

Broiler flocks: Before slaughter at farm

Bacteriological method: ISO 6579:2002

Broiler flocks: At slaughter (flock based approach)

Bacteriological method: ISO 6579:2002

Vaccination policy

Broiler flocks

no vaccination for salmonella is carried out in Malta.

Other preventive measures than vaccination in place

Broiler flocks

In accordance to Commission Regulation 1177/2007, no antimicrobial treatment is permitted as a control measure in the case of salmonella infection in poultry flocks.

Control program/mechanisms

The control program/strategies in place

Broiler flocks

Broiler flocks positive to S.typhimurium and S.enteritidis are culled. The A Salmonella National control programme according to Council Regulation 2160/2003 and Commission Regulation 646/2007 is being implemented.

Recent actions taken to control the zoonoses

Broiler flocks positive to s.typhimurium and s.enteritidis are destroyed.

Measures in case of the positive findings or single cases

Broiler flocks: Before slaughter at farm

Broiler flocks positive to s.typhimurium and s.enteritidis are destroyed.

Notification system in place

the National Veterinary Laboratory under the competent authority is responsible for notification of results to the Animal Health section and The Food Safety section which fall under the same CA for necessary action.

National evaluation of the recent situation, the trends and sources of infection

The Salmonella control programme has begun to be implemented in mid-2009, therefore it is a bit early to evaluate the situation . On what information is available the prevalence for all types of Salmonella infection is 31%.

C. Salmonella spp. in Gallus Gallus - flocks of laying hens

Monitoring system

Sampling strategy

Laying hens flocks

The competent authority samples all layer flocks irrespective of the capacity as official controls but also currently those on behalf of the business operators.

Frequency of the sampling

Laying hens: Rearing period

__18__ weeks prior to moving

Laying hens: Production period

Every __15__ weeks

Type of specimen taken

Laying hens: Rearing period

Environmental sample: samples of faeces and dust

Laying hens: Production period

Environmental sample: samples of faeces and dust

Methods of sampling (description of sampling techniques)

Laying hens: Rearing period

Two faecal samples and two dust samples are collected from each house on a holding. Faecal samples are collected from the bottom tiers of the manure belt.

Laying hens: Production period

The same samples are taken as described above.

Diagnostic/analytical methods used

Laying hens: Rearing period

Bacteriological method: ISO 6579:2002

Laying hens: Production period

Bacteriological method: ISO 6579:2002

Vaccination policy

Laying hens flocks

Layers are not vaccinated locally.

Other preventive measures than vaccination in place

Laying hens flocks

No antibiotic treatment is permitted as a means of control in case of salmonella infection in flocks.

Control program/mechanisms

The control program/strategies in place

Laying hens flocks

There is the Salmonella National control Programme that has been drawn up in accordance the Council Regulation 2160/2003 and Commission Regulation 1168/2006 and Control strategies followed are as

commission regulation 1177/2006.

Recent actions taken to control the zoonoses

Salmonella enteritidis and Salmonella typhimurium positive flocks have restriction measures implemented. The eggs are considered not fit for human consumption and are destroyed or heat-treated.

Measures in case of the positive findings or single cases

Laying hens flocks

Salmonella enteritidis and Salmonella typhimurium positive flocks have restriction measures implemented. The eggs are considered not fit for human consumption and are destroyed or heat-treated.

Notification system in place

All samples are brought into the National Veterinary Laboratory which carries out the analyses. Typing of positive isolates is carried out by the National reference laboratory. The National Veterinary Laboratory which falls under the competent authority is then responsible for communicating the information to the animal health section and food safety section which fall under the same authority for any further action if necessary. Action is only taken for Salmonella enteritidis and salmonella typhimurium positive cases.

National evaluation of the recent situation, the trends and sources of infection

The Salmonella Control Programme commenced in mid-2009. Based on the current data 42% of the holdings are positive for all types of Salmonella. it is too early in the programme to evaluate the trend.

Table Salmonella in breeding flocks of Gallus gallus

| | No of flocks under control programme | Source of information | Sampling strategy | Sampler | Sample type | Sample Origin | Target Verification | Sampling unit | Units tested | Total units positive for Salmonella | S. Enteritidis |
|--|--------------------------------------|-----------------------|-------------------|--------------------------------|-------------------|------------------------------|---------------------|---------------|--------------|-------------------------------------|----------------|
| Gallus gallus (fowl) - breeding flocks, unspecified - adult - Control and eradication programmes | | | Census | Official and industry sampling | | | yes | | | | |
| | S. Hadar | S. Infantis | S. Typhimurium | S. Virchow | S. 1,4,[5],12:i:- | Salmonella spp., unspecified | | | | | |
| Gallus gallus (fowl) - breeding flocks, unspecified - adult - Control and eradication programmes | | | | | | | | | | | |

Table Salmonella in other poultry

| | No of flocks under control programme | Source of information | Sampling strategy | Sampler | Sample type | Sample Origin | Target Verification | Sampling unit | Units tested | Total units positive for Salmonella | S. Enteritidis |
|---|--------------------------------------|-----------------------|--------------------|--------------------------------|--|---------------|---------------------|---------------|--------------|-------------------------------------|----------------|
| Gallus gallus (fowl) - laying hens - day-old chicks - Control and eradication programmes | 102 | | Objective sampling | Official sampling | environmental sample > hatchery basket liner | | yes | Batch | 3 | 0 | 0 |
| Gallus gallus (fowl) - laying hens - during rearing period - Control and eradication programmes | | | Objective sampling | Official and industry sampling | animal sample > faeces | | | Flock | | | |
| Gallus gallus (fowl) - laying hens - adult - at farm - Control and eradication programmes | | | Objective sampling | Official and industry sampling | animal sample > faeces | | yes | Flock | 102 | 9 | 7 |
| Gallus gallus (fowl) - broilers - before slaughter - at farm - Control and eradication programmes | 567 | | Objective sampling | Official and industry sampling | environmental sample > boot swabs | | yes | Flock | 561 | 4 | 3 |
| Turkeys - breeding flocks, unspecified - adult - at farm - Control and eradication programmes | | | Census | Official and industry sampling | | | yes | | | | |
| Turkeys - fattening flocks - before slaughter - at farm - Control and eradication programmes | | | Census | Official and industry sampling | | | yes | | | | |

| | S. Typhimurium | S. 1,4,[5],12:i:- | Salmonella spp., unspecified |
|---|-------------------|-------------------|------------------------------------|
| Gallus gallus (fowl) - laying hens - day-old chicks - Control and eradication programmes | 0 | 0 | 0 |
| Gallus gallus (fowl) - laying hens - during rearing period - Control and eradication programmes | | | |
| Gallus gallus (fowl) - laying hens - adult - at farm - Control and eradication programmes | 2 | 0 | 0 |

Table Salmonella in other poultry

| | S. Typhimurium | S. 1,4,[5],12:i:- | Salmonella spp., unspecified |
|---|-------------------|-------------------|------------------------------------|
| Gallus gallus (fowl) - broilers - before slaughter - at farm - Control and eradication programmes | 1 | 0 | 0 |
| Turkeys - breeding flocks, unspecified - adult - at farm - Control and eradication programmes | | | |
| Turkeys - fattening flocks - before slaughter - at farm - Control and eradication programmes | | | |

Footnote:

Data for breeding flocks of Gallus Gallus,turkey breeding flocks and turkey fattening flocks are not reported as there are no units or farms rearing or producing these catagories of poultry.

The laying hens day old chicks were reported as batch as these were tested prior to them being distributed to the respective flocks/holdings.

2.1.5 Salmonella in feedingstuffs

Table Salmonella in compound feedingstuffs

| | Source of information | Sampling strategy | Sampler | Sample type | Sample Origin | Sampling unit | Sample weight | Units tested | Total units positive for Salmonella | S. Enteritidis | S. Typhimurium |
|--|------------------------------|--------------------|-------------------|-------------|---------------|---------------|---------------|--------------|-------------------------------------|----------------|----------------|
| Compound feedingstuffs for poultry (non specified) - process control - at feed mill - Surveillance | | Objective sampling | Official sampling | feed sample | | Batch | | 8 | 0 | 0 | 0 |
| | Salmonella spp., unspecified | | | | | | | | | | |
| Compound feedingstuffs for poultry (non specified) - process control - at feed mill - Surveillance | 0 | | | | | | | | | | |

2.1.6 Antimicrobial resistance in Salmonella isolates

Table Antimicrobial susceptibility testing of Salmonella in humans

| Salmonella Isolates out of a monitoring program (yes/no) Number of isolates available in the laboratory | S. Enteritidis | | S. Typhimurium | | Salmonella spp. | | S. Bochum | | S. Concord | | S. Elomrane | | S. Hadar | | S. Haifa | | S. Infantis | | S. Kedougou | | S. Kentucky | | S. Livingstone | | S. London | |
|---|----------------|----|----------------|----|-----------------|----|-----------|---|------------|---|-------------|---|----------|---|----------|---|-------------|---|-------------|---|-------------|---|----------------|---|-----------|---|
| | no | | no | | no | | no | | no | | no | | no | | no | | no | | no | | no | | no | | no | |
| | 50 | | 24 | | 20 | | 1 | | 2 | | 1 | | 1 | | 4 | | 5 | | 1 | | 4 | | 2 | | 1 | |
| | N | n | N | n | N | n | N | n | N | n | N | n | N | n | N | n | N | n | N | n | N | n | N | n | N | n |
| Antimicrobials: | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Aminoglycosides - Gentamicin | 50 | 28 | 24 | 14 | 20 | 10 | 1 | 1 | 2 | 2 | 1 | 1 | 1 | 1 | 4 | 4 | 5 | 5 | 1 | 1 | 4 | 1 | 2 | 1 | 1 | 1 |
| Fluoroquinolones - Ciprofloxacin | 50 | 6 | 24 | 2 | 20 | 5 | 1 | 0 | 2 | 0 | 1 | 0 | 1 | 1 | 4 | 0 | 5 | 0 | 1 | 0 | 4 | 1 | 2 | 0 | 1 | 0 |
| Penicillins - Ampicillin | 50 | 8 | 24 | 16 | 20 | 7 | 1 | 0 | | | 1 | 0 | 1 | 1 | 4 | 3 | 5 | 2 | 1 | 0 | 4 | 0 | 2 | 0 | 1 | 0 |
| Trimethoprim + Sulfonamides | 50 | 6 | 24 | 2 | 20 | 5 | 1 | 0 | 2 | 1 | 1 | 0 | 1 | 1 | 4 | 1 | 5 | 0 | 1 | 0 | 4 | 0 | 2 | 0 | 1 | 0 |
| Fully sensitive | 50 | 31 | 24 | 18 | 20 | 12 | | | | | | | | | 4 | 0 | 5 | 4 | 1 | 1 | 4 | 1 | 2 | 1 | 1 | 1 |
| Resistant to 1 antimicrobial | 50 | 25 | 24 | 6 | 20 | 5 | 1 | 1 | | | 1 | 1 | | | 4 | 1 | 5 | 1 | 1 | 1 | 4 | 0 | 2 | 1 | 1 | 1 |
| Resistant to 2 antimicrobials | 0 | 0 | 24 | 10 | 20 | 3 | | | 2 | 1 | | | | | 4 | 2 | 5 | 3 | | | 2 | 1 | | | | |
| Resistant to 3 antimicrobials | 50 | 1 | | | | | | | 2 | 1 | | | | | 4 | 1 | | | | | | | | | | |
| Resistant to 4 antimicrobials | | | 24 | 2 | 20 | 4 | | | | | | | 1 | 1 | | | | | | | | | | | | |
| Resistant to >4 antimicrobials | 50 | 1 | | | | | | | | | | | | | | | | | | | | | | | | |

| Salmonella | S. Mkamba | | S. Newport | | S. Typhi | | S. Virchow | | S. enterica subsp. enterica | | S. enterica subsp. salamae | | Salmonella spp., unspecified | |
|----------------------------------|---|---|------------|---|----------|---|------------|---|-----------------------------|---|----------------------------|---|------------------------------|---|
| | Isolates out of a monitoring program (yes/no) | | | | | | | | | | | | | |
| | no | | no | | no | | no | | no | | no | | no | |
| | 1 | | 2 | | 2 | | 4 | | 3 | | 1 | | 3 | |
| Antimicrobials: | N | n | N | n | N | n | N | n | N | n | N | n | N | n |
| Aminoglycosides - Gentamicin | 1 | 1 | 2 | 1 | 2 | 1 | 4 | 4 | 3 | 0 | 1 | 1 | 3 | 3 |
| Fluoroquinolones - Ciprofloxacin | 1 | 0 | 2 | 0 | 2 | 0 | 4 | 4 | 3 | 0 | 1 | 0 | 3 | 1 |

Table Antimicrobial susceptibility testing of Salmonella in humans

| Salmonella | S. Mkamba | | S. Newport | | S. Typhi | | S. Virchow | | S. enterica subsp. enterica | | S. enterica subsp. salamae | | Salmonella spp., unspecified | |
|---|-----------|---|------------|---|----------|---|------------|---|--------------------------------|---|-------------------------------|---|------------------------------------|---|
| Isolates out of a monitoring program (yes/no) | no | | no | | no | | no | | no | | no | | no | |
| Number of isolates available in the laboratory | 1 | | 2 | | 2 | | 4 | | 3 | | 1 | | 3 | |
| Antimicrobials: | N | n | N | n | N | n | N | n | N | n | N | n | N | n |
| Penicillins - Ampicillin | 1 | 1 | 2 | 0 | 2 | 0 | 4 | 4 | 3 | 0 | 1 | 0 | 3 | 1 |
| Trimethoprim + Sulfonamides | 1 | 1 | 2 | 0 | 2 | 1 | 4 | 4 | 3 | 0 | 1 | 0 | 3 | 1 |
| Fully sensitive | 1 | 1 | 2 | 1 | 2 | 1 | 4 | 4 | 3 | 0 | 1 | 1 | 3 | 0 |
| Resistant to 1 antimicrobial | | | 2 | 1 | 2 | 1 | | | | | 1 | 1 | 3 | 2 |
| Resistant to 2 antimicrobials | | | | | | | | | | | | | | |
| Resistant to 3 antimicrobials | 1 | 1 | | | | | | | | | | | | |
| Resistant to 4 antimicrobials | | | | | | | 4 | 4 | | | | | 3 | 1 |
| Resistant to >4 antimicrobials | | | | | | | | | | | | | | |

Table Cut-off values for antibiotic resistance testing of Salmonella in Animals

| Test Method Used | Standard methods used for testing |
|------------------|-----------------------------------|
| | |

| | | | Concentration (microg/ml) | Zone diameter (mm) |
|------------------|-----------------|----------|---------------------------|--------------------|
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 2 | |
| | Streptomycin | | 32 | |
| Amphenicols | Chloramphenicol | | 16 | |
| Cephalosporins | Cefotaxime | | 0.5 | |
| Fluoroquinolones | Ciprofloxacin | | 0.06 | |
| Penicillins | Ampicillin | | 4 | |
| Quinolones | Nalidixic acid | | 16 | |
| Sulfonamides | Sulfonamides | | 256 | |
| Tetracyclines | Tetracycline | | 8 | |
| Trimethoprim | Trimethoprim | | 2 | |

Table Cut-off values for antibiotic resistance testing of Salmonella in Feed

| Test Method Used | | Standard methods used for testing | | |
|------------------|-----------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 2 | |
| | Streptomycin | | 32 | |
| Amphenicols | Chloramphenicol | | 16 | |
| Cephalosporins | Cefotaxime | | 0.5 | |
| Fluoroquinolones | Ciprofloxacin | | 0.06 | |
| Penicillins | Ampicillin | | 4 | |
| Quinolones | Nalidixic acid | | 16 | |
| Sulfonamides | Sulfonamides | | 256 | |
| Tetracyclines | Tetracycline | | 8 | |
| Trimethoprim | Trimethoprim | | 2 | |

| Test Method Used | Standard methods used for testing |
|------------------|-----------------------------------|
| | |

Table Cut-off values for antibiotic resistance testing of Salmonella in Humans

| Test Method Used | | Standard methods used for testing | | |
|-----------------------------|-----------------------------|-----------------------------------|---------------------------|--------------------|
| | | Eucast | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 4 | |
| Fluoroquinolones | Ciprofloxacin | | 1 | |
| Trimethoprim + Sulfonamides | Trimethoprim + Sulfonamides | | 80 | |

2.2 CAMPYLOBACTERIOSIS

2.2.1 General evaluation of the national situation

2.2.2 Campylobacteriosis in humans

A. Thermophilic Campylobacter in humans

Reporting system in place for the human cases

Statutorily notifiable.

Case definition

ECDC definition holds

Diagnostic/analytical methods used

Culture and serology

Notification system in place

Yes. Physician and laboratory surveillance system.

History of the disease and/or infection in the country

The rates of campylobacteriosis have been increasing in the past few years.

National evaluation of the recent situation, the trends and sources of infection

sources of infection are probably poultry meats. National campaigns and food safety weeks being held annually to educate the public.

Relevance as zoonotic disease

A relevant zoonotic disease in view of the numbers involved, morbidity, expenses and possible long term complications.

Table Campylobacter in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. | Autochthon cases | Autochthon Inc. | Imported cases | Imported Inc. | Unknown status |
|---------------------------------|-------|------------|------------------|-----------------|----------------|---------------|----------------|
| Campylobacter | 216 | 51.7 | 216 | 51.7 | 4 | .9 | 0 |
| C. coli | 33 | 8 | 33 | 8 | 1 | 0.2 | |
| C. jejuni | 103 | 24.7 | 103 | 24.7 | 3 | 0.7 | |
| C. upsaliensis | 0 | 0 | 0 | 0 | 0 | 0 | |
| Campylobacter spp., unspecified | 80 | 19 | 80 | 19 | 0 | 0 | |

Table Campylobacter in humans - Age distribution

| Age distribution | C. coli | | | C. jejuni | | | Campylobacter spp., unspecified | | |
|--------------------|---------|----|----|-----------|----|----|---------------------------------|----|----|
| | All | M | F | All | M | F | All | M | F |
| <1 year | 1 | 0 | 1 | 4 | 2 | 2 | 6 | 4 | 2 |
| 1 to 4 years | 8 | 5 | 3 | 28 | 18 | 10 | 7 | 3 | 4 |
| 5 to 14 years | 4 | 1 | 3 | 18 | 10 | 8 | 14 | 10 | 4 |
| 15 to 24 years | 5 | 0 | 5 | 13 | 4 | 9 | 5 | 2 | 3 |
| 25 to 44 years | 6 | 5 | 1 | 17 | 8 | 9 | 16 | 10 | 6 |
| 45 to 64 years | 6 | 3 | 3 | 4 | 3 | 1 | 10 | 5 | 5 |
| 65 years and older | 4 | 2 | 2 | 20 | 9 | 11 | 21 | 11 | 10 |
| Age unknown | 0 | 0 | 0 | 2 | 2 | 0 | 1 | 1 | 0 |
| Total : | 34 | 16 | 18 | 106 | 56 | 50 | 80 | 46 | 34 |

Table Campylobacter in humans - Seasonal distribution

| Seasonal Distribution Months | C. coli | C. jejuni | C. upsaliensi s | Campylobacter spp., unspecified |
|---------------------------------|---------|-----------|-----------------------|------------------------------------|
| | Cases | Cases | Cases | Cases |
| January | 5 | 8 | 0 | 0 |
| February | 2 | 5 | 0 | 3 |
| March | 1 | 5 | 0 | 12 |
| April | 0 | 14 | 0 | 18 |
| May | 3 | 16 | 0 | 6 |
| June | 2 | 10 | 0 | 4 |
| July | 6 | 4 | 0 | 5 |
| August | 0 | 8 | 0 | 5 |
| September | 2 | 5 | 0 | 7 |
| October | 3 | 8 | 0 | 4 |
| November | 3 | 11 | 0 | 6 |
| December | 7 | 12 | 0 | 10 |
| Total : | 34 | 106 | 0 | 80 |

2.2.3 Antimicrobial resistance in Campylobacter isolates

Table Antimicrobial susceptibility testing of Campylobacter in humans

| Campylobacter | Campylobacter spp., unspecified | | C. coli | | C. jejuni - C. jejuni subsp. jejuni | |
|--|---------------------------------|----|---------|----|-------------------------------------|----|
| Isolates out of a monitoring program (yes/no) | no | | no | | no | |
| Number of isolates available in the laboratory | 80 | | 34 | | 106 | |
| Antimicrobials: | N | n | N | n | N | n |
| Fluoroquinolones - Ciprofloxacin | 80 | 70 | 34 | 22 | 106 | 76 |
| Macrolides - Erythromycin | 80 | 65 | 34 | 27 | 106 | 24 |
| Fully sensitive | 80 | 75 | 34 | 22 | 106 | 76 |
| Resistant to 1 antimicrobial | 80 | 7 | 34 | 15 | 106 | 52 |
| Resistant to 2 antimicrobials | 80 | 64 | 34 | 7 | 106 | 24 |

Table Cut-off values used for antimicrobial susceptibility testing of Campylobacter in Humans

| Test Method Used | | Standard methods used for testing | | |
|------------------|--|-----------------------------------|--|--|
| E-test | | NCCLS/CLSI | | |

| | | | Concentration (microg/ml) | Zone diameter (mm) |
|------------------|---------------|----------|---------------------------|--------------------|
| | | Standard | Resistant > | Resistant <= |
| Fluoroquinolones | Ciprofloxacin | | 1 | |
| Macrolides | Erythromycin | | 4 | |

Table Cut-off values used for antimicrobial susceptibility testing of C. coli in Animals

| Test Method Used | | Standard methods used for testing | | |
|------------------|---------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 2 | |
| | Streptomycin | | 4 | |
| Fluoroquinolones | Ciprofloxacin | | 1 | |
| Macrolides | Erythromycin | | 16 | |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of C. coli in Feed

| Test Method Used | | Standard methods used for testing | | |
|------------------|---------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| Aminoglycosides | Gentamicin | | Concentration (microg/ml) | Zone diameter (mm) |
| | Streptomycin | Standard | Resistant > | Resistant <= |
| Fluoroquinolones | Ciprofloxacin | | 2 | |
| | | | 4 | |
| Macrolides | Erythromycin | | 1 | |
| Tetracyclines | | | 16 | |
| | Tetracycline | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of C. coli in Food

| Test Method Used | | Standard methods used for testing | | |
|------------------|---------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| Aminoglycosides | Gentamicin | | Concentration (microg/ml) | Zone diameter (mm) |
| | Streptomycin | Standard | Resistant > | Resistant <= |
| Fluoroquinolones | Ciprofloxacin | | 2 | |
| | | | 4 | |
| Macrolides | Erythromycin | | 1 | |
| Tetracyclines | | | 16 | |
| | Tetracycline | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of *C. jejuni* in Animals

| Test Method Used | | Standard methods used for testing | | |
|------------------|---------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| Aminoglycosides | Gentamicin | | Concentration (microg/ml) | Zone diameter (mm) |
| | Streptomycin | Standard | Resistant > | Resistant <= |
| Fluoroquinolones | Ciprofloxacin | | 1 | |
| | | | 2 | |
| Macrolides | Erythromycin | | 1 | |
| Tetracyclines | | | 4 | |
| | Tetracycline | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of *C. jejuni* in Feed

| Test Method Used | | Standard methods used for testing | | |
|------------------|---------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 1 | |
| | Streptomycin | | 2 | |
| Fluoroquinolones | Ciprofloxacin | | 1 | |
| Macrolides | Erythromycin | | 4 | |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of *C. jejuni* in Food

| Test Method Used | | Standard methods used for testing | | |
|------------------|---------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 1 | |
| | Streptomycin | | 2 | |
| Fluoroquinolones | Ciprofloxacin | | 1 | |
| Macrolides | Erythromycin | | 4 | |
| Tetracyclines | Tetracycline | | 2 | |

2.3 LISTERIOSIS

2.3.1 General evaluation of the national situation

A. Listeriosis general evaluation

History of the disease and/or infection in the country

No cases of human disease in recent years.

2.3.2 Listeriosis in humans

A. Listeriosis in humans

Reporting system in place for the human cases

A notifiable disease.

Case definition

ECDC definition holds.

Diagnostic/analytical methods used

Culture

Notification system in place

Yes

History of the disease and/or infection in the country

No notified cases of listeriosis in recent years.

2.4 E. COLI INFECTIONS

2.4.1 General evaluation of the national situation

2.5 TUBERCULOSIS, MYCOBACTERIAL DISEASES

2.5.1 General evaluation of the national situation

A. Tuberculosis general evaluation

History of the disease and/or infection in the country

In a recent study (1995-2005) the elderly group of locally born persons are mostly affected with an incidence of 10.6/100,000 persons-years. Since 2003 we have seen a rise in TB corresponding to a wave of illegal immigrants from the sub-sahharin regions of Africa.

National evaluation of the recent situation, the trends and sources of infection

The trends of TB in immigrants has been steady with a number of active and latent cases being treated.

Relevance of the findings in animals, feedingstuffs and foodstuffs to human cases (as a source of infection)

Immigrant cases are mostly imported ones. Few might be locally acquired possibly because of close proximity of living quarters in detention centers

Recent actions taken to control the zoonoses

routine screening of all immigrants and DOTS treatment.

2.5.2 Tuberculosis, mycobacterial diseases in humans

A. Tuberculosis due to Mycobacterium bovis in humans

Reporting system in place for the human cases

Statutory Notification from Laboratories, Physicians.

Diagnostic/analytical methods used

Culture

Notification system in place

Statutory Notification from Laboratories, Physicians.

History of the disease and/or infection in the country

very rare in humans

National evaluation of the recent situation, the trends and sources of infection

Last human case in 2006. Case acquired from contact with cattle. No further cases since.

Table Mycobacterium in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. | Autochtho n cases | Autochtho n Inc. | Imported cases | Imported Inc. |
|--------------------------------|-------|------------|----------------------|---------------------|-------------------|------------------|
| Mycobacterium | 25 | 5.99 | 5 | 1.2 | 20 | 0 |
| M. bovis | 0 | 0 | 0 | 0 | 0 | 0 |
| M. tuberculosis | 22 | 5.27 | 3 | 0.72 | 19 | |
| Reactivation of previous cases | 3 | 0.72 | 2 | 0.48 | 1 | |

Footnote:

Most of our TB cases occur in migrants from Sub-Saharan Africa (17 of 19 cases) and therefore imported.

The imported incidence of TB in Migrants (legal and illegal migrants) cannot be calculated as the denominator is unknown.

Table Mycobacterium in humans - Age distribution

| Age distribution | M. bovis | | | M. tuberculosis - complex | | |
|--------------------|----------|---|---|---------------------------|----|---|
| | All | M | F | All | M | F |
| <1 year | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 0 | 0 | 0 | 6 | 4 | 2 |
| 25 to 44 years | 0 | 0 | 0 | 11 | 8 | 3 |
| 45 to 64 years | 0 | 0 | 0 | 2 | 1 | 1 |
| 65 years and older | 0 | 0 | 0 | 3 | 2 | 1 |
| Age unknown | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : | 0 | 0 | 0 | 22 | 15 | 7 |

2.5.3 Mycobacterium in animals

A. Mycobacterium bovis in bovine animals

Status as officially free of bovine tuberculosis during the reporting year

Additional information

The last confirmed positive case was in 2001.

Monitoring system

Diagnostic/analytical methods used

On farm skin test carried out twice yearly according to standard protocols.

Measures in case of the positive findings or single cases

The positive animal is slaughtered and tissue samples taken for microbiology tests. If microbiology is negative the farm is considered as negative.

Notification system in place

The Veterinary officer in charge of animal health is responsible of informing the farmer. The farmer is then advised to make arrangements for the slaughter of the animal and sampling for Microbiology is carried out.

Table Bovine tuberculosis in countries and regions that do not receive Community co-financing for eradication programmes

If present, the row "Total -1" refers to analogous data of the previous year.

| Region | Total number of existing bovine | | Officially free herds | | Infected herds | | Routine tuberculin testing | | Number of tuberculin tests carried out before the introduction into the herds (Annex A(I)(2)(c) third indent (1) of Directive 64/432/EEC) | Number of animals with suspicious lesions of tuberculosis examined and submitted to histopathological and bacteriological | Number of animals detected positive in bacteriological examination |
|-----------------------|---------------------------------|---------|-----------------------|-----|-----------------|---|---|--------------------------|---|---|--|
| | Herds | Animals | Number of herds | % | Number of herds | % | Interval between routine tuberculin tests | Number of animals tested | | | |
| Malta | 125 | 13912 | 125 | 100 | 0 | 0 | once a year | 13765 | 0 | 0 | 0 |
| Total : ¹⁾ | 125 | 13912 | 125 | 100 | 0 | 0 | N.A. | 13765 | 0 | 0 | 0 |

Comments:

¹⁾ N.A.

Footnote:

The total number of holdings with bovine animals is 324, 125 of these holdings are strictly milk producing holdings thus while there are 324 holdings under the Brucella programme there are only 125 holdings under the Tuberculosis programme.

2.6 BRUCELLOSIS

2.6.1 General evaluation of the national situation

A. Brucellosis general evaluation

History of the disease and/or infection in the country

The last case of brucellosis in Malta occurred in 1998. This followed a short epidemic related to goat's milk in 1995.

National evaluation of the recent situation, the trends and sources of infection

Strict control of animal herds have eliminated human cases since 1998.

Relevance of the findings in animals, feedingstuffs and foodstuffs to human cases (as a source of infection)

n reported findings in foodstuffs.

Recent actions taken to control the zoonoses

Routine surveillance from the veterinary end.

2.6.2 Brucellosis in humans

A. Brucellosis in humans

Reporting system in place for the human cases

Statutory notification obliging laboratories and physicians to notify cases.

Case definition

Clinical picture with demonstration of specific antibody response, demonstration by immunofluorescence of Isolation of Brucella.

Diagnostic/analytical methods used

Serology, Immunofluorescence and isolation from clinical specimens.

Notification system in place

Statutory notification by labs and physicians.

History of the disease and/or infection in the country

As above.

National evaluation of the recent situation, the trends and sources of infection

NO human cases since 1998.

Relevance as zoonotic disease

There is the hope that in the future Malta might be declared Brucella free

Table Brucella in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. | Autochthon cases | Autochthon Inc. | Imported cases | Imported Inc. |
|-------------------------------|-------|------------|------------------|-----------------|----------------|---------------|
| Brucella | 0 | 0 | 0 | 0 | 0 | 0 |
| B. abortus | 0 | 0 | 0 | 0 | 0 | 0 |
| B. melitensis | 0 | 0 | 0 | 0 | 0 | 0 |
| B. suis | 0 | 0 | 0 | 0 | 0 | 0 |
| Occupational cases | 0 | 0 | 0 | 0 | 0 | 0 |

Table Brucella in humans - Age distribution

| Age distribution | B. abortus | | | B. melitensis | | | Brucella spp., unspecified | | |
|--------------------|------------|---|---|---------------|---|---|----------------------------|---|---|
| | All | M | F | All | M | F | All | M | F |
| <1 year | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 to 44 years | 0 | 0 | 0 | 0 | | 0 | 0 | 0 | 0 |
| 45 to 64 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 65 years and older | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Age unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2.6.3 Brucella in animals

A. Brucella abortus in bovine animals

Status as officially free of bovine brucellosis during the reporting year

The entire country free

Brucella abortus has never been recorded in bovines in Malta. The last case confirmed in Bovines was B.melitensis in 1996.

Monitoring system

Sampling strategy

All animals over 6 months

Frequency of the sampling

Blood twice Yearly, Milk bulk tank 3 times a year.

Type of specimen taken

Other: blood, milk or tissue

Methods of sampling (description of sampling techniques)

Blood sampling on farms is carried out twice yearly for screening with RBT/ELISA. Bulk milk tank samples are taken 3 times a year to screen by MRT.

Diagnostic/analytical methods used

Rose Bengal Test, ELISA, Milk Ring Test, Complement Fixation Test, Microbiological analysis of lymph node samples from CFT positive animals.

Vaccination policy

No vaccination for brucellosis is carried out in Malta.

Other preventive measures than vaccination in place

Animal movement is controlled and only authorized by the CA and based on the Health Status of the farm.

Control program/mechanisms

The control program/strategies in place

Animals that test positive to a complement fixation test are slaughtered and tissue samples lifted. These samples are then subject to microbiological analysis.

Measures in case of the positive findings or single cases

The positive animal is slaughtered and tissue samples lifted. These samples are then sent for microbiology. All the animals on the farm are retested at an interval of 1 month from the slaughter of the CFT positive animal and again after 2 months. If the microbiology test is negative the farm is considered negative.

Notification system in place

The National Veterinary Laboratory sends the results of screening to the National Livestock database. In the case of a positive screening result the Veterinary officer responsible for Animal Health is informed by email of the positive result. Farmers are sent a copy of both the negative and positive result (RBT,MRT,ELISA,Microbiology). If the microbiology test is negative the farm is considered negative.

B. Brucella melitensis in goats

Status as officially free of caprine brucellosis during the reporting year

The entire country free

Brucella abortus has never been recorded in caprines . The last case confirmed was B.melitensis in 1996.

Monitoring system

Sampling strategy

All animals over 6 months are tested twice yearly.

Frequency of the sampling

All animals over 6 months are tested twice yearly.

Type of specimen taken

Blood

Methods of sampling (description of sampling techniques)

Blood samples are taken from all animals over 6 months.

Diagnostic/analytical methods used

Rose Bengal Test, Complement Fixation Test, ELISA and Microbiology

Vaccination policy

No Vaccination is practiced in the Maltese Islands

Other preventive measures than vaccination in place

Strict control of animal movement

Measures in case of the positive findings or single cases

Any animal positive for RBT is then re tested using CFT and ELISA. Positive CFT/ELISA animals are slaughtered and organs are submitted for Microbiological investigation.

Table Ovine or Caprine Brucellosis in countries and regions that do not receive Community co-financing for eradication programme

If present, the row "Total -1" refers to analogous data of the previous year.

| Region | Total number of existing | | Officially free herds | | Infected herds | | Surveillance | | | Investigations of suspect cases | | | | |
|-----------------------|--------------------------|---------|-----------------------|-----|-----------------|---|------------------------|--------------------------|--------------------------|---|--|--|--|---------------------------|
| | Herds | Animals | Number of herds | % | Number of herds | % | Number of herds tested | Number of animals tested | Number of infected herds | Number of animals tested with serological blood tests | Number of animals positive serologically | Number of animals examined microbiologically | Number of animals positive microbiologically | Number of suspended herds |
| Malta | 2188 | 16984 | 2188 | 100 | 0 | 0 | 1406 | 14932 | 0 | 14932 | 0 | 0 | 0 | 0 |
| Total : ¹⁾ | 2188 | 16984 | 2188 | 100 | 0 | 0 | 1406 | 14932 | 0 | 14932 | 0 | 0 | 0 | 0 |

Comments:

¹⁾ N.A.

Table Bovine brucellosis in countries and regions that do not receive Community co-financing for eradication programme

If present, the row "Total -1" refers to analogous data of the previous year.

| Region | Total number of existing bovine | | Officially free herds | | Infected herds | | Surveillance | | | | | | Investigations of suspect cases | | | | | | | | |
|-----------------------|---------------------------------|---------|-----------------------|------|-----------------|---|-------------------------------|--------------------------|--------------------------|-------------------------------|-----------------------------------|--------------------------|---|--|---|---|---------------------------|----------------------------|-----|--|--|
| | | | | | | | Serological tests | | | Examination of bulk milk | | | Information about | | | Epidemiological investigation | | | | | |
| | Herds | Animals | Number of herds | % | Number of herds | % | Number of bovine herds tested | Number of animals tested | Number of infected herds | Number of bovine herds tested | Number of animals or pools tested | Number of infected herds | Number of notified abortions whatever cause | Number of isolations of Brucella infection | Number of abortions due to Brucella abortus | Number of animals tested with serological blood tests | Number of suspended herds | Number of positive animals | | Number of animals examined microbiologically | Number of animals positive microbiologically |
| | | | | | | | | | | | | | | | | | | Sero logically | BST | | |
| Malta | 324 | 15545 | 254 | 78.4 | 0 | 0 | 254 | 13126 | 0 | 125 | 239 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : ¹⁾ | 324 | 15545 | 254 | 78.4 | 0 | 0 | 254 | 13126 | 0 | 125 | 239 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Comments:

¹⁾ N.A.

2.7 YERSINIOSIS

2.7.1 General evaluation of the national situation

A. Yersinia enterocolitica general evaluation

History of the disease and/or infection in the country

No cases in recent years.

2.7.2 Yersiniosis in humans

A. Yersiniosis in humans

Reporting system in place for the human cases

Statutorily notifiable infection.

Case definition

Clinical criteria and Isolation of pathogenic bacteria in a clinical specimen.

Diagnostic/analytical methods used

Testing in accredited to foreign labs.

Notification system in place

Yes

History of the disease and/or infection in the country

No recent history of disease locally.

2.8 TRICHINELLOSIS

2.8.1 General evaluation of the national situation

A. Trichinellosis general evaluation

History of the disease and/or infection in the country

There are no recorded human or animal cases.

National evaluation of the recent situation, the trends and sources of infection

The possibility of swine being infected with trichinella are very remote. There are no wild animals in Malta that may support the cycle in wildlife. All Pigs are reared indoors. There are no backyard farms and no freerange pigs

Recent actions taken to control the zoonoses

There is no wildlife on Malta. Trichinella testing in the past was carried out using the trichinoscope method. End of 2009 the digestive method was being adopted. Full analysis of all horse/swine slaughtered commenced beginning 2010.

2.8.2 Trichinellosis in humans

A. Trichinellosis in humans

Reporting system in place for the human cases

Statutorily notifiable disease.

Case definition

EDC definition holds.

Diagnostic/analytical methods used

Bopsy of skeletal muscle and serological testing.

Notification system in place

From physicians and laboratories.

History of the disease and/or infection in the country

No recorded human cases.

Table Trichinella in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. | Autochthon cases | Autochthon Inc. | Imported cases | Imported Inc. |
|-------------------------------|-------|------------|------------------|-----------------|----------------|---------------|
| Trichinella | 0 | 0 | 0 | 0 | 0 | 0 |
| Trichinella spp., unspecified | 0 | 0 | 0 | 0 | 0 | 0 |

Table Trichinella in humans - Age distribution

| Age distribution | Trichinella spp., unspecified | | |
|--------------------|-------------------------------|---|---|
| | All | M | F |
| <1 year | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | |
| 15 to 24 years | 0 | 0 | 0 |
| 25 to 44 years | 0 | 0 | 0 |
| 45 to 64 years | 0 | 0 | 0 |
| 65 years and older | 0 | 0 | 0 |
| Age unknown | 0 | 0 | 0 |
| Total : | 0 | 0 | 0 |

2.8.3 Trichinella in animals

A. Trichinella in horses

Monitoring system

Sampling strategy

All Horses slaughtered for human consumption are sampled

Frequency of the sampling

All Horses slaughtered for human consumption are sampled

Type of specimen taken

Muscle

Diagnostic/analytical methods used

Digestive Method as per council regulation 2075/2005

Table Trichinella in animals

| | Source of information | Sampling strategy | Sampler | Sample type | Sample Origin | Sampling unit | Units tested | Total units positive for Trichinella | T. spiralis | Trichinella spp., unspecified |
|--|--------------------------------|--------------------|-------------------|------------------------------|---------------|---------------|--------------|--------------------------------------|-------------|-------------------------------|
| Pigs - fattening pigs - raised under controlled housing conditions - at slaughterhouse - Surveillance | National Veterinary Laboratory | Objective sampling | Official sampling | animal sample > organ/tissue | | Single | 81089 | 0 | 0 | 0 |
| Pigs - breeding animals - raised under controlled housing conditions - sows and boars - at slaughterhouse - Surveillance | National Veterinary Laboratory | Objective sampling | Official sampling | animal sample > organ/tissue | | Single | 2321 | 0 | 0 | 0 |
| Solipeds, domestic - horses - at slaughterhouse - Surveillance | National Veterinary Laboratory | Objective sampling | Official sampling | animal sample > organ/tissue | | Single | 76 | 0 | 0 | 0 |

2.9 ECHINOCOCCOSIS

2.9.1 General evaluation of the national situation

A. Echinococcus spp. general evaluation

History of the disease and/or infection in the country

No notified cases in the past few years

2.9.2 Echinococcosis in humans

A. Echinococcus spp. in humans

Reporting system in place for the human cases

Statutorily notifiable disease.

Case definition

ECDC definition holds

Notification system in place

Routine surveillance

Additional information

No cases have been reported.

Table Echinococcus in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. | Autochthon cases | Autochthon Inc. | Imported cases | Imported Inc. |
|--------------------------------|-------|------------|------------------|-----------------|----------------|---------------|
| Echinococcus | 0 | 0 | 0 | 0 | 0 | 0 |
| E. granulosus | 0 | 0 | 0 | 0 | 0 | 0 |
| E. multilocularis | 0 | 0 | 0 | 0 | 0 | 0 |
| Echinococcus spp., unspecified | 0 | 0 | 0 | 0 | 0 | 0 |

Table Echinococcus in humans - Age distribution

| Age distribution | E. granulosus | | | E. multilocularis | | | Echinococcus spp., unspecified | | |
|--------------------|---------------|---|---|-------------------|---|---|--------------------------------|---|---|
| | All | M | F | All | M | F | All | M | F |
| <1 year | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 to 24 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 to 44 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 45 to 64 years | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 65 years and older | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Age unknown | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total : | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

2.10 TOXOPLASMOSIS

2.10.1 General evaluation of the national situation

2.10.2 Toxoplasmosis in humans

A. Toxoplasmosis in humans

Reporting system in place for the human cases

Statutorily notifiable disease.

Case definition

ECDC definition holds.

Diagnostic/analytical methods used

Serological methods.

Notification system in place

Yes

History of the disease and/or infection in the country

No notified cases.

Table Toxoplasma in humans - Species/serotype distribution

| Species/serotype Distribution | Cases | Cases Inc. |
|-------------------------------|-------|------------|
| Toxoplasma | 0 | 0 |
| Toxoplasma spp., unspecified | 0 | 0 |
| Congenital cases | 0 | 0 |

Table Toxoplasma in humans - Age distribution

| Age distribution | Toxoplasma spp., unspecified | | |
|--------------------|------------------------------|---|---|
| | All | M | F |
| <1 year | 0 | 0 | 0 |
| 1 to 4 years | 0 | 0 | 0 |
| 5 to 14 years | 0 | 0 | 0 |
| 15 to 24 years | 0 | 0 | 0 |
| 25 to 44 years | 0 | 0 | 0 |
| 45 to 64 years | 0 | 0 | 0 |
| 65 years and older | 0 | 0 | 0 |
| Age unknown | 0 | 0 | 0 |
| Total : | 0 | 0 | 0 |

2.10.3 Toxoplasma in animals

Table Toxoplasma in animals

| | Source of information | Sampling strategy | Sampler | Sample type | Sample Origin | Analytical Method | Sampling unit | Units tested | Total units positive for Toxoplasma | T. gondii | Toxoplasma spp., unspecified |
|------------------------------|--------------------------------|--------------------|-------------------|-----------------------|---------------|-------------------|---------------|--------------|-------------------------------------|-----------|------------------------------|
| Sheep - at farm - Monitoring | National Veterinary Laboratory | Selective sampling | Official sampling | animal sample > blood | | ELISA | Animal | 400 | 311 | 311 | 0 |
| Goats - at farm - Monitoring | National Veterinary Laboratory | Selective sampling | Official sampling | animal sample > blood | | ELISA | Animal | 409 | 273 | 273 | 0 |

2.11 RABIES

2.11.1 General evaluation of the national situation

A. Rabies general evaluation

History of the disease and/or infection in the country

Last reported rabies case at the beginning of the 20th century!

A notifiable infection

Relevance of the findings in animals, feedingstuffs and foodstuffs to human cases (as a source of infection)

No known animal cases.

2.11.2 Rabies in humans

A. Rabies in humans

Reporting system in place for the human cases

Rabies is still a notifiable disease in Malta.

Case definition

ECDC definition holds.

2.12 STAPHYLOCOCCUS INFECTION

2.12.1 General evaluation of the national situation

2.13 Q-FEVER

2.13.1 General evaluation of the national situation

A. *Coxiella burnetii* (Q-fever) general evaluation

History of the disease and/or infection in the country

No reported cases in recent years.

3. INFORMATION ON SPECIFIC INDICATORS OF ANTIMICROBIAL RESISTANCE

3.1 ESCHERICHIA COLI, NON-PATHOGENIC

3.1.1 General evaluation of the national situation

A. Escherichia coli general evaluation

History of the disease and/or infection in the country

E. coli O157 disease is not common with a handful of cases each year.

National evaluation of the recent situation, the trends and sources of infection

No recent trends noticed.

Recent actions taken to control the zoonoses

Educational campaigns, stricter control at food production premises.

3.1.2 Antimicrobial resistance in Escherichia coli, non-pathogenic

Table Cut-off values used for antimicrobial susceptibility testing of Escherichia coli, non-pathogenic in Animals

| Test Method Used | | Standard methods used for testing | | |
|------------------|-----------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 2 | |
| | Streptomycin | | 16 | |
| Amphenicols | Chloramphenicol | | 16 | |
| Cephalosporins | Cefotaxime | | 0.25 | |
| Fluoroquinolones | Ciprofloxacin | | 0.03 | |
| Penicillins | Ampicillin | | 8 | |
| Quinolones | Nalidixic acid | | 16 | |
| Sulfonamides | Sulfonamides | | 256 | |
| Tetracyclines | Tetracycline | | 8 | |
| Trimethoprim | Trimethoprim | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of Escherichia coli, non-pathogenic in Animals

Table Cut-off values used for antimicrobial susceptibility testing of Escherichia coli, non-pathogenic in Feed

| Test Method Used | | Standard methods used for testing | | |
|------------------|-----------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 2 | |
| | Streptomycin | | 16 | |
| Amphenicols | Chloramphenicol | | 16 | |
| Cephalosporins | Cefotaxime | | 0.25 | |
| Fluoroquinolones | Ciprofloxacin | | 0.03 | |
| Penicillins | Ampicillin | | 8 | |
| Quinolones | Nalidixic acid | | 16 | |
| Sulfonamides | Sulfonamides | | 256 | |
| Tetracyclines | Tetracycline | | 8 | |
| Trimethoprim | Trimethoprim | | 2 | |

Table Cut-off values used for antimicrobial susceptibility testing of Escherichia coli, non-pathogenic in Food

| Test Method Used | | Standard methods used for testing | | |
|------------------|-----------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 2 | |
| | Streptomycin | | 16 | |
| Amphenicols | Chloramphenicol | | 16 | |
| Cephalosporins | Cefotaxime | | 0.25 | |
| Fluoroquinolones | Ciprofloxacin | | 0.03 | |
| Penicillins | Ampicillin | | 8 | |
| Quinolones | Nalidixic acid | | 16 | |
| Sulfonamides | Sulfonamides | | 256 | |
| Tetracyclines | Tetracycline | | 8 | |
| Trimethoprim | Trimethoprim | | 2 | |

3.2 ENTEROCOCCUS, NON-PATHOGENIC

3.2.1 General evaluation of the national situation

3.2.2 Antimicrobial resistance in Enterococcus, non-pathogenic isolates

Table Cut-off values for antibiotic resistance of *E. faecalis* in Animals

| Test Method Used | | Standard methods used for testing | | |
|---|---------------------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 32 | |
| | Streptomycin | | 512 | |
| Amphenicols | Chloramphenicol | | 32 | |
| Glycopeptides (Cyclic peptides, Polypeptides) | Vancomycin | | 4 | |
| Macrolides | Erythromycin | | 4 | |
| Oxazolidines | Linezolid | | 4 | |
| Penicillins | Ampicillin | | 4 | |
| Streptogramins | Quinupristin/Dalfopristin | | 32 | |

Table Cut-off values for antibiotic resistance of *E. faecalis* in Animals

| | | | Concentration (microg/ml) | Zone diameter (mm) |
|---------------|--------------|----------|---------------------------|--------------------|
| | | Standard | Resistant > | Resistant <= |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values for antibiotic resistance of E. faecalis in Feed

| Test Method Used | | Standard methods used for testing | | |
|---|---------------------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 32 | |
| | Streptomycin | | 512 | |
| Amphenicols | Chloramphenicol | | 32 | |
| Glycopeptides (Cyclic peptides, Polypeptides) | Vancomycin | | 4 | |
| Macrolides | Erythromycin | | 4 | |
| Oxazolidines | Linezolid | | 4 | |
| Penicillins | Ampicillin | | 4 | |
| Streptogramins | Quinupristin/Dalfopristin | | 32 | |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values for antibiotic resistance of E. faecalis in Food

| Test Method Used | | Standard methods used for testing | | |
|---|---------------------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 32 | |
| | Streptomycin | | 512 | |
| Amphenicols | Chloramphenicol | | 32 | |
| Glycopeptides (Cyclic peptides, Polypeptides) | Vancomycin | | 4 | |
| Macrolides | Erythromycin | | 4 | |
| Oxazolidines | Linezolid | | 4 | |
| Penicillins | Ampicillin | | 4 | |
| Streptogramins | Quinupristin/Dalfopristin | | 32 | |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values for antibiotic resistance of E. faecium in Animals

| Test Method Used | | Standard methods used for testing | | |
|---|---------------------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 32 | |
| | Streptomycin | | 128 | |
| Amphenicols | Chloramphenicol | | 32 | |
| Glycopeptides (Cyclic peptides, Polypeptides) | Vancomycin | | 4 | |
| Macrolides | Erythromycin | | 4 | |
| Oxazolidines | Linezolid | | 4 | |
| Penicillins | Ampicillin | | 4 | |
| Streptogramins | Quinupristin/Dalfopristin | | 1 | |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values for antibiotic resistance of E. faecium in Feed

| Test Method Used | | Standard methods used for testing | | |
|---|---------------------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 32 | |
| | Streptomycin | | 128 | |
| Amphenicols | Chloramphenicol | | 32 | |
| Glycopeptides (Cyclic peptides, Polypeptides) | Vancomycin | | 4 | |
| Macrolides | Erythromycin | | 4 | |
| Oxazolidines | Linezolid | | 4 | |
| Penicillins | Ampicillin | | 4 | |
| Streptogramins | Quinupristin/Dalfopristin | | 1 | |
| Tetracyclines | Tetracycline | | 2 | |

Table Cut-off values for antibiotic resistance of E. faecium in Food

| Test Method Used | | Standard methods used for testing | | |
|---|---------------------------|-----------------------------------|---------------------------|--------------------|
| | | | | |
| | | | Concentration (microg/ml) | Zone diameter (mm) |
| | | Standard | Resistant > | Resistant <= |
| Aminoglycosides | Gentamicin | | 32 | |
| | Streptomycin | | 128 | |
| Amphenicols | Chloramphenicol | | 32 | |
| Glycopeptides (Cyclic peptides, Polypeptides) | Vancomycin | | 4 | |
| Macrolides | Erythromycin | | 4 | |
| Oxazolidines | Linezolid | | 4 | |
| Penicillins | Ampicillin | | 4 | |
| Streptogramins | Quinupristin/Dalfopristin | | 1 | |
| Tetracyclines | Tetracycline | | 2 | |

4. INFORMATION ON SPECIFIC MICROBIOLOGICAL AGENTS

4.1 ENTEROBACTER SAKAZAKII

4.1.1 General evaluation of the national situation

A. Enterobacter sakazakii general evaluation

History of the disease and/or infection in the country

No known history of cases in Malta.

National evaluation of the recent situation, the trends and sources of infection

No cases of disease identified in Malta

Recent actions taken to control the hazard

Surveillance being done on infant milk formulations at the laboratory end.

Suggestions to the Community for the actions to be taken

None so far

4.1.2 Cronobacter in foodstuffs

A. Enterobacter sakazakii in foodstuffs

Monitoring system

Type of specimen taken

Milk

Methods of sampling (description of sampling techniques)

EC regulation 2073/2005 on microbiological criteria on food stuffs where in 30 samples of 10 grms all samples must be negative.

Definition of positive finding

Detection of Enterobacter sakazakii in any of the samples.

Diagnostic/analytical methods used

ISO/ts: 22964 /2006

Preventive measures in place

Monitoring programme.

4.2 HISTAMINE

4.2.1 General evaluation of the national situation

A. Histamine General evaluation

History of the disease and/or infection in the country

Malta has few reported cases of histamine poisoning in humans every year. It may be under reported.

Relevance of the findings in animals, feedingstuffs and foodstuffs to human cases (as a source of infection)

Most cases occur via canned tuna or from locally caught dolphin fish which were temperature mis-treated at some stage of food preparation.

4.3 STAPHYLOCOCCAL ENTEROTOXINS

4.3.1 General evaluation of the national situation

A. Staphylococcal enterotoxins general evaluation

History of the disease and/or infection in the country

Not common.

National evaluation of the recent situation, the trends and sources of infection

There are no particular trends being observed

Recent actions taken to control the hazard

Educational campaigns on food safety and hygiene undergoing.

5. FOODBORNE

Foodborne outbreaks are incidences of two or more human cases of the same disease or infection where the cases are linked or are probably linked to the same food source. Situation, in which the observed human cases exceed the expected number of cases and where a same food source is suspected, is also indicative of a foodborne outbreak.

A. Foodborne outbreaks

System in place for identification, epidemiological investigations and reporting of foodborne outbreaks

Statutory notification system involving all physicians and laboratories. Epidemiological investigation done by epidemiologists at the Health promotion and Disease Prevention Directorate, with field investigations performed by the Environmental Health Directorate of the Regulatory directorate of the Health Department.

Description of the types of outbreaks covered by the reporting:

All suspected types of food-borne outbreaks reported, however not all outbreaks are reported. All notified outbreaks are classified according to origin.

National evaluation of the reported outbreaks in the country:

Trends in numbers of outbreaks and numbers of human cases involved

There has been no major trend over the last decade in the number of outbreaks. Minor changes from year to year reflect random variability as the country is small.

Relevance of the different type of places of food production and preparation in outbreaks

All types of food-borne outbreaks are more common in households (30-45%). The next most common source are restaurants.

Evaluation of the severity and clinical picture of the human cases

No specific trends have been noticed.

Table Foodborne Outbreaks: summarised data

| | Weak evidence or no vehicle outbreaks | | | | Strong evidence Number of Outbreaks | Total number of outbreaks |
|---|---------------------------------------|-------------|--------------|---------|-------------------------------------|---------------------------|
| | Number of outbreaks | Human cases | Hospitalized | Deaths | | |
| Salmonella - S. Typhimurium | 1 | 2 | 1 | 0 | 0 | 1 |
| Salmonella - S. Enteritidis | 4 | 24 | 3 | 0 | 0 | 4 |
| Salmonella - Other serovars | 1 | 2 | 1 | 0 | 0 | 1 |
| Campylobacter | 12 | 25 | 5 | 0 | 0 | 12 |
| Listeria - Listeria monocytogenes | 0 | unknown | unknown | unknown | 0 | 0 |
| Listeria - Other Listeria | 0 | unknown | unknown | unknown | 0 | 0 |
| Yersinia | 0 | unknown | unknown | unknown | 0 | 0 |
| Escherichia coli, pathogenic - Verotoxigenic E. coli (VTEC) | 0 | unknown | unknown | unknown | 0 | 0 |
| Bacillus - B. cereus | 0 | unknown | unknown | unknown | 0 | 0 |
| Bacillus - Other Bacillus | 0 | unknown | unknown | unknown | 0 | 0 |
| Staphylococcal enterotoxins | 0 | unknown | unknown | unknown | 0 | 0 |
| Clostridium - Cl. botulinum | 0 | unknown | unknown | unknown | 0 | 0 |
| Clostridium - Cl. perfringens | 0 | unknown | unknown | unknown | 0 | 0 |

| | Weak evidence or no vehicle outbreaks | | | | Strong evidence Number of Outbreaks | Total number of outbreaks |
|---|---------------------------------------|-------------|--------------|---------|-------------------------------------|---------------------------|
| | Number of outbreaks | Human cases | Hospitalized | Deaths | | |
| Clostridium - Other Clostridia | 0 | unknown | unknown | unknown | 0 | 0 |
| Other Bacterial agents - Brucella | 0 | unknown | unknown | unknown | 0 | 0 |
| Other Bacterial agents - Shigella | 2 | 15 | 2 | 0 | 0 | 2 |
| Other Bacterial agents - Other Bacterial agents | 0 | unknown | unknown | unknown | 0 | 0 |
| Parasites - Trichinella | 0 | unknown | unknown | unknown | 0 | 0 |
| Parasites - Giardia | 0 | unknown | unknown | unknown | 0 | 0 |
| Parasites - Cryptosporidium | 0 | unknown | unknown | unknown | 0 | 0 |
| Parasites - Anisakis | 0 | unknown | unknown | unknown | 0 | 0 |
| Parasites - Other Parasites | 0 | unknown | unknown | unknown | 0 | 0 |
| Viruses - Norovirus | 3 | 46 | 0 | 0 | 0 | 3 |
| Viruses - Hepatitis viruses | 0 | unknown | unknown | unknown | 0 | 0 |
| Viruses - Other Viruses | 0 | unknown | unknown | unknown | 0 | 0 |
| Other agents - Histamine | 4 | 11 | 0 | 0 | 0 | 4 |
| Other agents - Marine biotoxins | 0 | unknown | unknown | unknown | 0 | 0 |
| Other agents - Other Agents | 0 | unknown | unknown | unknown | 0 | 0 |

Unknown agent

| Weak evidence or no vehicle outbreaks | | | | Strong evidence Number of Outbreaks | Total number of outbreaks |
|---------------------------------------|-------------|--------------|--------|-------------------------------------|---------------------------|
| Number of outbreaks | Human cases | Hospitalized | Deaths | | |
| 33 | 124 | 4 | 0 | 0 | 33 |