



European Food Safety Authority

**Workshop on Acrylamide Formation in Food
17 November 2003, Brussels**

Report of the Workshop*

1. SUMMARY

The meeting demonstrated that since April 2002 international research on acrylamide in food has progressed rapidly, however there are still many questions that need to be answered. The interest in the acrylamide issue is apparent from the response to this meeting: 60 participants from the food processing and catering industry, governmental bodies and academia from Europe, Canada and the USA attended the workshop. A full list of attendees is given in Annex II.

During the course of discussions several areas were identified to further investigate the formation of acrylamide in food. The results of this work may then be used to inform methods of reducing acrylamide levels. It was highlighted that there is a need to develop a better understanding of the fundamental chemistry of acrylamide, how it is formed, what the rate-limiting steps of formation are and how formation can be reduced. The main challenge for researchers is to understand how acrylamide is formed and how to influence the mechanism of formation, in order to reduce acrylamide levels whilst retaining the food's nutritional and organoleptic properties, and not adversely affecting other food safety parameters.

Areas identified as needing further research included:

- The role of water and the physical state of the food matrix.
- Time and temperature processing parameters.
- The influence of ammonia/ammonium bicarbonate and other CO₂-releasing agents on acrylamide levels.
- Chemical or enzymatic reduction of acrylamide e.g. using asparaginase to reduce asparagine levels.
- Developing profiles of acrylamide precursors in raw materials and examining the effects of natural variation (e.g. crop variety, weather, season and agronomic practices) on these. This information could be used to advise farmers, concerning husbandry and crop storage to potentially minimise the acrylamide precursors, or could be transferred into a plant-breeding programme.
- Optimising storage conditions of potatoes to obtain a balance between acrylamide production potential and organoleptic properties.

* This report reviews the current knowledge and ideas for future work, as described by researchers who attended the workshop. The comments provided are the opinions of the workshop attendees and do not necessarily reflect the views of the European Food Safety Authority (EFSA).

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

2. BACKGROUND

In 2002 Stockholm University and the Swedish National Food Authority announced the results of research which showed that unexpectedly high levels of acrylamide were generated in a wide range of starch-rich foods cooked at high temperatures¹. Further research has since confirmed these findings.

In 1991 the EC's SCF (see Annex I for a glossary of terms and abbreviations) evaluated acrylamide as a monomer in food contact materials and concluded that acrylamide is a genotoxic carcinogen. In light of the Swedish findings the SCF considered the implications of acrylamide formation in food in July 2002 and recommended that levels of acrylamide in food should be as low as reasonably achievable (ALARA)². At this stage the SCF could only offer general advice due to the lack of detailed knowledge on scientific issues relevant to the risk management. On an international level, the WHO/FAO held a joint consultation to assess the full extent of the public health risk from acrylamide in food³. Both this consultation and the SCF recommended that further research be conducted on acrylamide in food, to cover topics such as formation, toxicity and possible means of reduction.

In response to the recommendations made by the SCF and the WHO/FAO consultation, the VWA and the FSA, in association with the CIAA, held the 28 March 2003 Workshop "Research on Acrylamide in Food"⁴. The outcomes and recommendations of the meeting were detailed in a "White Paper on Acrylamide" and presented to the EFSA Advisory Forum⁴. Following EFSA's consideration of the White Paper, a "Consultation Paper" was presented to the EFSA Advisory Forum, requesting that EFSA take the lead role in two of the six recommendations made at the meeting: to formulate a research plan to fill the gaps in current knowledge and co-ordinate the implementation of individual projects, and secondly to create a mechanism for information exchange and discussion, envisaged as a forum/task force for technical level discussion⁵.

As a result of these activities and willingness from stakeholders the 17 November 2003 EFSA Workshop on Acrylamide Formation in Food was organised on EFSA's behalf by the FSA and VWA, with assistance from CIAA and the EC.

3. WORKSHOP OVERVIEW

The workshop provided an opportunity for those conducting research on acrylamide formation in food to engage in technical level discussions on current research activity, both within and beyond the EU. The aim of the workshop was to stimulate an exchange of ideas and findings so as to enhance the current understanding of the mechanisms of formation and facilitate an effective and sustainable means of reducing levels of acrylamide in foods and to thereby protect consumer health. This workshop was specifically a technical level discussion looking at research being conducted on formation of acrylamide in foods. The analytical methodology, toxicology, and methods of reducing acrylamide were not within the scope of this workshop.

The Chair, Dr Herman Koëter, opened the workshop with an overview of the aims and expected outcomes, and an introduction to EFSA's role in co-ordinating acrylamide activities. As an introduction to the workshop and to stimulate ideas for

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

the discussion group sessions, presentations were given by representatives of the EFSA, EC, CIAA and US JIFSAN. The workshop agenda can be found in Annex III.

Following this introductory plenary session, participants were assigned to two discussion groups made up of individuals from a mix of research backgrounds and expertise. Each group participated in informal discussions, covering a range of issues from the intricacies of the chemistry of formation to the effects of processing techniques on acrylamide formation.

4. REPORT OF DISCUSSION GROUPS

The outcomes of the discussion groups are summarised below. It must be taken into consideration that in certain circumstances further work may be required to confirm results described and this is highlighted in the text. As well as looking at how understanding the formation can be used to reduce acrylamide levels, the effects of changing food production methods should also be considered, for example the effects on food safety parameters, nutrient content and organoleptic qualities.

Much of the research carried out to date has been completed in model systems. The use of model systems is effective in testing the importance of physical factors; however, when using model systems it is important to remember that they are simplifications of complex food systems and therefore caution should be taken when relating findings to real foods.

4.1. Chemistry of Formation

To fully understand the acrylamide concentrations observed in certain foods, it is important to consider both the formation and elimination reactions of acrylamide. Evidence so far shows that acrylamide is eliminated by reaction with numerous food constituents, such as inherent nucleophiles (e.g. amines and sulfhydryls) and that thermal degradation products of sugars and starch may also play a role. Rapid elimination reactions may provide an explanation for the low levels of acrylamide found in particular food types, for example meat⁶. Also, above certain temperatures, acrylamide levels have been shown to decrease, i.e. where the rate of elimination exceeds the rate of formation.

Although several mechanisms for the formation of acrylamide have been proposed, the precise mechanism(s) or pathway(s) for acrylamide formation in food have not yet been fully elucidated. Further knowledge on the precise mechanistic pathway is required and the rate-limiting steps need to be identified with the help of kinetic studies.

4.1.1 Formation of Acrylamide via Asparagine and the Maillard Reaction

Several research groups^{6,7,8,9,10,11} have independently proposed that the main pathway to acrylamide formation in foodstuffs is by the reaction between asparagine and a carbonyl source (e.g. reducing sugars). In the Maillard reaction, acrylamide may be formed by at least two competing pathways starting from a common precursor, the Schiff base. This could lead via classical Strecker degradation to the Strecker aldehyde of asparagine⁷, or the Schiff base could undergo rapid decarboxylation and subsequent beta-elimination to form acrylamide¹².

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

Glucose and fructose are the important reactants so far identified. In the case of potatoes, fructose is approximately twice as efficient at promoting acrylamide formation than glucose, and 15 times more than lactose^{6,13}. The amino acid asparagine is the source of the nitrogen in acrylamide, this observation has been corroborated by nitrogen-15 labelling studies¹⁴. Other amino acids present in food will react with the available reducing sugars, competing with asparagine, which consequently may lead to a reduction in the levels of acrylamide. The quantity of acrylamide formed is largely dependent on the availability of the reducing sugar and asparagine. The rate of formation can be approximated by a bimolecular reaction between fructose/glucose and asparagine^{13,15}. However, other subsequent steps are also likely to contribute to determining the rate. In cereal products the rate-limiting factor appears to be the availability of asparagine, whereas in potato products the availability of reducing sugar is the determining factor. In model reactions between amino acids and glucose it has been demonstrated that the yield of acrylamide is approximately 0.1%^{9,13}.

There were discussions about the role of sucrose as an alternative carbonyl source. The general opinion appeared to be that sucrose does not contribute to the formation of acrylamide, however, it may be important in supporting elimination of acrylamide in some foods, such as, gingerbread. Other carbonyls in food (some being intermediates in the Maillard pathway) may also react with asparagine to form acrylamide.

The formation of acrylamide appears to be strongly linked to the Maillard reaction and hence the favourable qualities of colour, texture and flavour that are associated with roast and fried foods. Acrylamide formation is therefore linked to some of the desirable organoleptic qualities in such foods.

4.1.2 Additional Acrylamide Formation Pathways

It was the general view of the participants that oxidation of lipids and starch did not significantly contribute to the formation of acrylamide in food. Acrylamide is generally formed in starch-rich foods, however, the influence of starch on acrylamide formation is thought to be limited to physical rather than chemical properties e.g. its effect on the dry matter content.

In model experiments, other pathways have been identified that may lead to the formation of acrylamide from reaction precursors, such as, acrolein, acrylic acid and other carbonyls^{16,17} (Table 1). In starch-rich foods it was agreed that asparagine is the main source of nitrogen in the acrylamide molecule. However, other potential sources have been identified, for example, when heating lipids in the absence of carbohydrate, ammonia is an alternative source of nitrogen in the acrylamide molecule, provided via the deamination of amino acids¹⁷. This work suggests that ammonia and acrolein could play an important role in acrylamide formation in lipid-rich foods. However, these alternative formation mechanisms have yet to be demonstrated in real food systems.

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

Table 1: Summary of known acrylamide formation mechanisms in model experiments

Acrylamide Formation Mechanism	Reference
<ul style="list-style-type: none"> Intramolecular cyclisation of Schiff base to oxazolidin-5-one and –CO₂, beta-elimination of the decarboxylated Amadori product to afford acrylamide 	Yaylayan et al. 2003 ¹²
<ul style="list-style-type: none"> Also non-classical Maillard compounds + Asparagine lead to Acrylamide formation. Thus, two mechanisms may be operating. Methionine (or other amino acids) may be active after –CO₂, –NH₂ to methional, behaves as an aldehyde + Asparagine ↔ Acrylamide Acrolein + NH₃ ↔ Acrylamide or, acrolein + Asparagine ↔ Acrylamide by the Maillard pathway Acrylic acid (from Asparagine or other degradation products) + NH₃ ↔ Acrylamide 	Friedman, M. J. 2003 ¹⁸
<ul style="list-style-type: none"> Acrylic acid + NH₃ ↔ Acrylamide (ammonia from thermal degradation of amino acids) Only approx. 5% of the yield compared to Asparagine 	Stadler et al. 2003 ¹⁶
<ul style="list-style-type: none"> Acrolein (from triolein) + Asparagine ↔ Acrylamide (88.6 µg/g) Acrolein + NH₃ ↔ Acrylamide (753 µg/g) Acrylic acid + NH₃ ↔ Acrylamide (190 000 µg/g) 	Yasuhara et al. 2003 ¹⁷
<ul style="list-style-type: none"> Non-oxidative mechanism proposed: acrylic acid from 2-propenal 	Vattem & Shetty, 2003 ¹⁹
<ul style="list-style-type: none"> Formation from a common precursor (decarboxylated Schiff base) by two routes: beta elimination to Acrylamide or 3-aminopropionamide and deamination ↔ Acrylamide 	Zyzak et al., 2003 ¹⁴
<ul style="list-style-type: none"> Heating polyacrylamide at 175°C (15, 30 min) in presence of transition metals 	Ahn et al., 2003 ²⁰

Many studies have investigated the formation and reduction of acrylamide, looking at the pH dependence, reduction by addition of enzymes in model systems e.g. asparaginase in dough and potato models²¹. Studies in model systems have also shown that the elimination of acrylamide can be accelerated by addition of heated (caramelised) sugars^{13,22}. Addition of sugars not only promotes acrylamide formation by providing a carbonyl source in the Maillard reaction but may also reduce acrylamide under certain conditions.

The addition of proteins to model systems has been investigated using fish²³, meat and chickpea proteins¹⁹ and has resulted in lower acrylamide levels. It has been demonstrated that coating potato slices with chickpea batter before frying results in a reduction in acrylamide levels¹⁹. The mechanism is still unknown, however suggestions include the chemical elimination of acrylamide or the formation of a thermal barrier. These observations may explain the low levels of acrylamide observed in high protein foods.

4.1.3 Gaps Highlighted and Potential Areas of Research in Acrylamide Formation

- Maillard Reaction:
 - Fundamental research on the mechanistic chemistry of acrylamide outside of the known food systems and the Maillard reaction.
 - What is the efficiency of the Maillard reaction (reducing sugars and carbonyls) to produce acrylamide? Why is the yield of acrylamide formation observed in model systems high in comparison to other Maillard reaction products (as observed in many foods such as roasted nuts, coffee and certain cereals), taking into consideration that elimination reactions commence already at temperatures above 120°C? What limits the acrylamide formation?

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

Dependent on the conditions, in many foods a significant proportion of free asparagine and sugars remain unreacted.

- How does the addition of heated sugars influence the final acrylamide levels?
- Work on potato model systems showed that acrylamide levels decreased initially at about 120°C⁶, this effect is more prominent in coffee that is roasted at temperatures above 200°C²⁴. It has been suggested that there is a reduction of acrylamide at high temperatures due to the lack of available reducing sugars. Further work is required to investigate whether there are other mechanisms contributing.
- Are there other precursors that have not yet been identified? Kinetic modelling will assist in determining the rate-limiting steps and thus provide a means of potentially controlling acrylamide formation.
- Amino Acid Mixtures:
 - The competitive effect of free amino acid mixtures must be investigated and linked to the free amino acid pool of the food matrix.
 - There is a need to map the total free amino acid and sugar profiles in crops and raw materials and how this influences acrylamide formation, this could also be extended to other specific amino acids such as lysine and glycine.
 - Kinetic modelling could be used to understand the role of asparagine versus the role of the total amino acids or other specific amino acids.

As a result of these discussions, a need was identified to better understand the fundamental chemistry of acrylamide formation, elucidate the fundamental steps of the reaction and then assess under real food processing conditions, which of the possible pathways are dominant. This understanding should then be translated to food systems and the chemistry that governs industrial food processing, domestic and catering food preparation practices, so that the benefits of the Maillard reaction can be realised without producing acrylamide.

4.2 Physical Parameters

The groups discussed the role of the food matrix and in particular starch and proteins involved in acrylamide formation. Acrylamide formation has been associated with starch-rich foods, however, it has not been shown thus far that starch itself is involved in the formation of acrylamide in food. It has been hypothesised that the starch matrix could entrap, in some manner, acrylamide precursors, intermediates or acrylamide itself, or that the dry matter content of the food effects formation. Another hypothesis is that starch provides an inert matrix, which does not promote elimination of acrylamide; therefore rates of formation are greater than elimination. It should also be considered that starch may have no correlation with acrylamide formation, but it is just that foods containing high levels of asparagine and reducing sugars are also starch-rich.

Temperature has an important role in the formation and elimination of acrylamide. It is well known that acrylamide forms in foods that are cooked at high temperatures, however, when temperatures are raised further an increase in acrylamide destruction is apparent^{6,7}. In cereal products the more often a product is heated the higher the destruction of acrylamide, and may be related to the secondary heating effects. It was recommended that further work should be carried out in order to support this hypothesis. In some cases higher baking temperatures are associated with less acrylamide, suggesting that there is a balance of simultaneous reactions forming and

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

destroying acrylamide¹³. There may be a critical temperature/time zone where acrylamide is formed at a greater rate than it is destroyed, at temperatures outside of this zone no/little acrylamide is present. In an attempt to reduce the length of time that foods are cooked at critical temperatures, the use of flash frying and rapid cooling could be used to reduce acrylamide formation. However, it has been suggested that these processes do not produce products with optimum organoleptic properties.

Using thermal imaging techniques, the distribution of temperature in food as it bakes has been studied for some types of cereal biscuit. The imaging shows graphically what is well known to experts in the energy efficiency of baking. The loss of water as the food dries during baking cools it and the bulk of the product is at a temperature very much lower than that of the oven. Temperature, time and moisture are key drivers of acrylamide formation, but the temperature of the food is not that of the oven and the moisture content is at a gradient from centre to crust. Small scale chemistry research needs to embrace the physics of heat and water transport in baking if results are to be meaningfully interpreted and transferred to a manufacturing scale. However, these products are not homogeneous and the observed results will therefore vary considerably.

It was agreed that the water content and the physical state of the food matrix could affect the mechanistic pathway for acrylamide formation. Water affects the chemical route as well as the mobility of the chemical constituents in polymeric matrices such as starch¹⁴, i.e. the transition from the rubbery to glassy state. A small increase in the final water content (from 1% to 2%) has been shown to reduce acrylamide formation in potato crisps²¹. In potato crisps it is thought that the low water content is one of the key contributors to high acrylamide levels¹³. Water management may be a key factor in controlling acrylamide in foods and warrants study in both industrial and domestic processing.

In order to determine how to reduce acrylamide levels effectively, it is important to know whether the formation of acrylamide is controlled by chemical or physical processes, and to gain further understanding of the acrylamide formation process and the Maillard reaction.

Several areas were identified which require further research:

- It has been suggested that acrylamide may only be present at high levels within a critical temperature zone, where the formation reaction is greater than elimination. Further work is required to determine whether altering processing techniques could avoid this critical temperature zone.
- Water content could be a key factor to controlling acrylamide levels, but little is still known about the effects of water content and the physical state of the food matrix on acrylamide formation.

4.3. Processing Techniques and Recipes

Processing techniques as well as the composition of the raw materials determine product characteristics (e.g. colour, flavour, and texture) and influence acrylamide levels in foodstuffs. Several researchers to date have shown that there is a relationship between product colour and acrylamide levels^{25,26}, however, there must be a colour/texture balance in order to have a product with acceptable organoleptic

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

properties. Conversely, other researchers believe that product colour cannot be relied on as a good indicator of acrylamide levels.

In experiments looking at the production of roasted potato products, it has been shown that reducing sugar levels of 1g/kg and 2g/kg in fresh weight potatoes resulted in acrylamide levels of approximately 500ppb and 1000ppb respectively²⁷. These researchers therefore concluded that potatoes with less than 1 g/kg reducing sugar are desired to retain low levels of acrylamide in roasted or fried potatoes. It has been seen that acrylamide levels can be significantly reduced by decreasing the reducing sugar content. However, because potato products are not uniform, both acrylamide and reducing sugar levels can vary considerably between products and therefore it is not possible to fix a maximum level of reducing sugars in fresh potatoes destined for cooked potato products.

In a Swiss study participants were asked to prepare French fries following preprepared instructions. This guidance detailed methods of preparing the fries to minimise acrylamide levels, without compromising product quality. 63 % of the participants produced fries with less than 100 ppb.

It was pointed out that processing techniques are much more complex in other sectors, such as the baking industry, where several unique and complex processes/recipe combinations are employed for individual product types. A much better understanding of the chemistry is required before significant changes can consistently be made.

Processing temperature alone is an inadequate parameter to compare different processing techniques. Heat flux allows the measurement of the energy transferred to the food from conductive, convective and radiative processes in an oven, and is therefore a more accurate representation of the processes occurring in the oven. The rapid cooling of low density, fried snack products following cooking can be demonstrated to lower acrylamide levels. In this instance rapid cooling prevents further cooking post frying and hence additional acrylamide generation.

Based on industrial baking trials and model experiments, it has been observed that the use of ammonium carbonate as a baking aid can increase acrylamide formation¹³. It is not obvious what role ammonia plays in acrylamide formation; when ammonium carbonate was added to a glucose–asparagine system (without a food matrix) no increase in acrylamide was observed. However, in some model experiments the addition of ammonium carbonate has resulted in an increased yield¹³. Reducing the amount of the raising agent ammonium bicarbonate was shown to reduce the acrylamide levels in plain flour matrices and the addition of lactic acid had a similar effect. Yung et al showed that the addition of citric acid to both French fries (1-2 %) and corn chips (0.1 – 0.2%) reduced acrylamide levels²⁸. However, citric acid accelerates the degradation of the frying oil that consequently leads to off-flavours and unacceptable quality of the products.

The role of ammonia needs to be explored and this was identified as a gap in the current knowledge. In the meantime it was felt that there is a need to investigate other CO₂-releasing agents that are compatible with food processing technology and will not compromise the desired organoleptic characteristics. The addition of SO₂ as an

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

antioxidant was discussed as an effective means of reducing acrylamide levels during processing, although its use is limited by its potential allergenic properties²⁹.

Other processing techniques which have been seen to produce a reduction in acrylamide levels, include the chemical or enzymatic reduction of asparagine, or the addition of amino acids other than asparagine. Addition of cysteine has been shown to be most effective in the reduction of acrylamide, although it produced an unpleasant off-flavour in the food. Blanching potatoes before cooking leaches out sugars from the potato and therefore reduces the acrylamide formation potential²⁶.

The affect of cooking oil has also been investigated, but due to the water-soluble nature of acrylamide it is not thought to be important; in a recent study it has been shown that acrylamide is not found in the oil following cooking. One question that remains is the effect of using old cooking oils on the level of acrylamide, and the potential contribution from the accumulation of particulates on the oil filters during frying. However, further work is required to establish how significant this effect would be.

Areas for further investigation:

- The role of ammonia in acrylamide formation.
- The consequences of high temperatures and the addition of sugars as elimination options.
- Process-related parameters (high temperature, high sugar) and different forms of cooking processes that result in high acrylamide levels.
- Feasibility of acrylamide reduction in coffee and other commodities.

For all heating processes the formation of acrylamide is related to the three physical factors, temperature, time and water content, and this may in turn affect the safety and organoleptic qualities of the food. Therefore when changing processing techniques to alter acrylamide formation, processors must also consider the expectations and safety of the consumer.

4.4. Agricultural Effects

Variation in raw materials and ingredients play an important role in acrylamide formation due to natural variation in levels of asparagine and reducing sugars. Reducing sugar levels in potatoes vary considerably with a number of factors including crop variety, seasonal changes, growing conditions (e.g. weather and type of soil) and storage conditions. In cereals it appears that variation in asparagine levels could be an important factor determining the acrylamide formation potential, although further data are required. Potato crops also show variation in reducing sugar and asparagine levels both between and within batches¹⁵, hence this is an important problem.

In both cereals and potatoes, it has been seen that varying the crops growing conditions will effect the variation in precursors, these differences are more significant in potatoes. It is thought that there is a higher homogeneity of precursors within a batch of grain than in potatoes, however, further work is needed to confirm these observations. The selection of crop varieties with lower reducing sugar levels provides a limited reduction in the levels of acrylamide and is currently being used in some areas of the cereal market. Millers blend all kinds of grain and use wide

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

varieties of flours, and also add minerals and other additives; this makes the selection of cereal products based on crop type more difficult. Asparagine levels can vary depending on the degree of milling, for example, dark flours can contain twice as much asparagine as white flour³⁰, however, in alternative experiments no significant difference has been observed.

Genetic modification of plant materials could provide a quick and effective reduction of reducing sugars and therefore the potential for acrylamide formation, for example, inhibiting the enzyme invertase to reduce the formation of reducing sugars in potatoes. This would avoid the lengthy process of crop breeding programmes, but due to other considerations it is not currently possible to use genetically engineered products.

The use of asparaginase in yeast during bread production was also discussed and is said to successfully reduce levels of acrylamide in bread, although the fermentation time is significantly increased. This is an important area for manufacturers of leavened dough products and further investigations are required to discover how it affects acrylamide formation.

Asparagine levels vary between varieties of cereals and little is known about what affects these levels. Potatoes produced using organic fertilisers have been shown to result in potatoes with slightly higher asparagine levels. Additionally the nitrogen content of the soil also appears to have a stronger link to the amino acid content in potatoes than it does in cereals, however, some experiments have shown that nitrogen fertilisers have no effect on the levels of asparagine^{6,15}.

The time that potatoes are harvested may also have a significant effect on the potential for acrylamide formation. The maturity stage of the potato tubers affects the reducing sugar levels. If two crops of the same variety, grown under the same conditions were harvested at different times, the more immature potatoes would have higher levels of reducing sugar. This relative difference in sugar levels will remain the same if the potatoes are kept under the same storage conditions³¹.

In order to develop a sustainable approach to reducing acrylamide levels in the long term, it was felt that information on the effect of the fundamental chemistry of the precursors on acrylamide levels (e.g. free amino acid and sugar profiles), must be translated into plant breeding programs. However, it was noted that the benefits from such programmes may take more than 10 years to be realised. The challenge will be to consistently achieve the desired levels of precursors crop after crop within seasonal and climatic variations.

Comparatively little is known about cereals and other commodities such as coffee and the effects different agricultural practices have on the levels of acrylamide in the final foods. Further work is required on different crop varieties, critical components and processing parameters effecting the formation of acrylamide in coffee. It is also necessary to identify other potential commodities that may produce high levels of acrylamide.

Further research is required to determine the free amino acid pool and reducing sugar content for targeted raw materials e.g. wheat, other cereals, coffee and cocoa, and the

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

associated natural variations within a single crop, within a season and over a number of years. For example what are the effects of seasons with low levels of precipitation on the levels of acrylamide formed in the final processed foodstuff?

Areas for further investigation for the development of a plant-breeding programme:

- Bio-agricultural research is needed to understand the metabolic profiles and develop fingerprints for the key precursors, and how this affects the composition of priority raw materials and hence acrylamide formation.
- Environmental affects on the composition of raw materials and how this affects acrylamide formation potential, for example, the use of fertilisers, climatic variation and natural variation, both between crops and from fluctuations between years and seasons.

4.5. Storage Conditions

Controlling storage conditions of potatoes could be a simple method for reducing acrylamide levels in cooked potato products³². It has been widely recognised that the storage conditions of potatoes can contribute significantly to the potential for acrylamide formation. By carrying out a systematic assessment of storage parameters, such as, temperature, humidity and light on factors such as levels of asparagine, reducing sugars and the associated product quality, this may provide information on reducing the potential for acrylamide formation. Storage of potatoes at temperatures below 8°C and storage over long periods of time have been shown to increase the concentration of reducing sugars and asparagine²¹, which consequently increases the potential for acrylamide formation. It was also suggested that asparagine is not effected by storage conditions, therefore further work is recommended to confirm these suggestions. Recommendations were made for research to assess the associated risk of increasing storage temperatures above 4°C, including the potential for disease and the effects of using sprout inhibitors on product quality and safety. It was thought that some of these factors could be overcome by selecting crops that were suited for particular storage conditions.

It was suggested that potatoes destined for industrial processing are currently kept under controlled storage conditions, whereas fresh potatoes to be sold for domestic use are stored at lower temperatures, therefore increasing the potential for high reducing sugar levels and therefore the formation of acrylamide in domestically prepared potato products.

It has been found that reconditioning potatoes by storing them at higher temperatures for 2-3 weeks will lower levels of reducing sugars and consequently the potential for acrylamide formation. When considering this it must be acknowledged that it generally takes at least 2 weeks from delivery to processing in the home, which could provide the necessary reconditioning. However, reconditioning for 2-3 weeks will not reduce reducing sugar levels to the original levels³².

As well as attempting to lower reducing sugar levels and therefore the potential for acrylamide formation, it may also be possible, by monitoring levels of reducing sugars in batches of raw potatoes, to recommend products suitable for particular processes.

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

It is currently thought that the storage conditions of cereals do not have a significant effect on the potential for acrylamide formation. Little is known about the effects on other commodities and this would therefore benefit from additional research.

During discussions the following areas were highlighted as requiring further research:

- The systematic assessment of storage parameters, such as, temperature, light and humidity on acrylamide precursor levels.
- Associated risk of changing storage conditions on product safety and quality, and the potential effects of using sprout inhibitors on potatoes.

4.7. Communication and Information Exchange

Continued use should be made of the systems already in place, for example, populating the EC JRC Monitoring Database on Acrylamide Levels³³ with data generated from national surveillance programmes and other analyses. Also keeping updated the Acrylamide Information Base of Research Activities in the EU³⁴ and the FAO/WHO Acrylamide Infonet³⁵.

The format for the discussion and exchange of new ideas was not discussed. Closer contact and interaction should be made with potato research institutes, e.g. The Federal Research Centre for Nutrition and Food, Germany (formerly The Federal Centre for Cereal Potato and Lipid Research), Wageningen UR or Agrotechnology and Food Innovations, Netherlands. A project on selection of cultivars and storage conditions to supply potatoes for roasting, baking and frying is being carried out by the Swiss Federal Research Station for Agroecology and Agriculture FAL, Reckenholz-Zurich.

Efforts should be made to exchange information on the formation of acrylamide through communication with the EC 6th Framework Programme HEATOX project³⁶, via its dissemination package. Also the University of Newcastle is carrying out a study on Low Input Farming as part of the EC 6th Framework Programme³⁶. This project aims to lead to new technologies and systems for organic and low input production systems, the research will include potatoes and cereals.

5. CONCLUSIONS

The key conclusion highlighted by many participants of the workshop was that further research was required to better understand the fundamental chemistry of acrylamide formation. It was noted by several participants that for a number of the relevant foods there is insufficient information available to be able to consistently lower the levels of acrylamide formed. There is a need to elucidate the fundamental steps of the reaction and then assess under real food processing conditions, which of the possible pathways are dominant. This understanding should then be translated to food systems and the chemistry that governs industrial food processing, domestic and catering food preparation practices; so that the benefits of the Maillard reaction can be realised without producing acrylamide and adversely affecting other safety/nutritional factors.

In addition to the research needs listed below other recommendations were:

- Continue the co-ordination of acrylamide activities via the Acrylamide Information Base of Research Activities in the EU and the FAO/WHO Acrylamide Infonet.

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

- Provide regular updates on research activities and the EC 6th Framework Programme Heatox Project.
- Liaise with the EC's DG Agriculture on relevant issues such as, potential plant breeding programmes, and collaborate with farmers, agriculture and research institutes.

During the discussions a number of areas were highlighted for further research:

5.1. Chemistry of Formation:

5.1.1. Maillard Reaction:

- Fundamental research on the mechanistic chemistry of acrylamide outside of the known food systems and the Maillard reaction.
- Investigate further the mechanism for acrylamide destruction at high temperatures.
- Identification of other potential acrylamide precursors.
- Investigate why low levels of acrylamide are observed in high protein foods.

5.1.2. Amino Acid Mixtures:

- Map the total free amino acid and sugar profiles in crops and raw materials and how this influences acrylamide formation.
- Kinetic modelling could be used to understand the role of asparagine versus the role of other amino acids.

5.2. Physical Parameters:

- Further work is required to determine whether altering processing techniques could avoid temperatures where the rate of acrylamide formation is greater than the rate of destruction.
- The effects of water content and the physical state of the food matrix on acrylamide formation.
- The physics of heat and water transport in baking.

5.3. Processing Techniques:

- The role of ammonia in acrylamide formation.
- Process-related parameters (high temperature, high sugar) and different forms of cooking processes that result in high acrylamide levels.
- The effect of using old cooking oils on the level of acrylamide, and the potential contribution from the accumulation of particulates on the oil filters during frying.

5.4. Agricultural Effects:

- Understand the metabolic profiles and develop fingerprints for the key precursors, and how this affects the composition of priority raw materials and hence acrylamide formation.
- Environmental affects on the composition of raw materials and how this affects acrylamide formation potential.

5.5. Storage Conditions:

- The systematic assessment of storage parameters, such as, temperature, light and humidity on acrylamide precursor levels.
- Associated risk of changing storage conditions on product safety and quality, including the use of sprout inhibitors on potatoes.

AF 06.04.2004 – 10 report on EFSA acrylamide workshop

Annex I Glossary of Terms and Abbreviations

Annex II Attendance List - EFSA Workshop on Acrylamide Formation in Food

Annex III Agenda - EFSA Workshop on Acrylamide Formation in Food

References

- ¹ Tareke, E., Rydberg, P., Karlsson, P., Eriksson, S. and Tornqvist, M. (2002) Analysis of Acrylamide, a carcinogen formed in heated foodstuffs. *Journal of Agricultural and Food Chemistry*, **51**(17), 4998-5006.
- ² European Commission Scientific Committee on Food (2002) Opinion of the Scientific Committee on Food on new findings regarding the presence of acrylamide in food http://europa.eu.int/comm/food/fs/sc/scf/out131_en.pdf
- ³ Health implications of acrylamide in food. Joint FAO/WHO consultation, Geneva, Switzerland, 25 - 27 June 2002 ISBN: 02 4 156218 8
http://www.who.int/foodsafety/publications/chem/acrylamide_june2002/en/
- ⁴ White Paper on Acrylamide, 3-4 July 2003 EFSA Advisory Forum Meetings, http://www.efsa.eu.int/af_meetings_en.html (UK Paper: Annex 3).
- ⁵ Acrylamide Consultation Paper, 3-4 July 2003 EFSA Advisory Forum Meetings, http://www.efsa.eu.int/af_meetings_en.html (UK Paper: Acrylamide Consultation Paper).
- ⁶ Biedermann, M.; Noti, A.; Biedermann-Brem, S.; Mozzetti, V. and Grob, K. (2002) Experiments on Acrylamide Formation and Possibilities to Decrease the Potential of Acrylamide Formation in Potatoes. *Mitteilungen aus Lebensmitteluntersuchung und Hygiene* **93**, 668-687
- ⁷ Mottram, D.S.; Wedzicha, B.L. & Dodson, A.T. (2002) Acrylamide is Formed in the Maillard Reaction. *Nature* **419**, 448-449
- ⁸ Sanders, R.A.; Zyzak, D.V.; Stojanovic, M.; Tallmadge, D.H.; Eberhart, B.L.; & Ewald, D.K. (2002) An LC/MS acrylamide method and its use in investigating the role of asparagine. *Acrylamide Symposium*, 116th Annual AOAC International Meeting, September 22-26, 2002, Los Angeles, California
- ⁹ Becalski, A.; Lau, B. P-Y.; Lewis, D. & Seaman S. W. (2003) Acrylamide in Foods: Occurrence, Sources and Modelling. *Journal of Agricultural and Food Chemistry* **51** (3), 802-808
- ¹⁰ Stadler, R.H.; Blank, I. Varga, N. Robert, F. Hau, J. Guy, PA. Robert, M-C. Riediker, S. (2002) Acrylamide from Maillard reaction products. *Nature* **419**, 449.
- ¹¹ Weißhaar, R. & Gutsche, B. (2002) Formation of Acrylamide in Heated Potato Products - Model Experiments Pointing to Asparagine as Precursor. *Deutsche Lebensmittel-Rundschau* **98**, 397-400
- ¹² Yaylayan, V.A.; Wnorowski, A., & Perez Locas C. (2003) Why asparagine needs carbohydrate to generate acrylamide. *Journal of Agricultural and Food Chemistry* **51**, 1753-1757
- ¹³ Biedermann, M. and Grob, K. (2003) Model studies on acrylamide formation in potato, wheat flour and corn starch; ways to reduce acrylamide contents in bakery ware. *Mitteilungen aus Lebensmitteluntersuchung und Hygiene* **94**, 406-422
- ¹⁴ Zyzak, DV. Sanders, RA. Stojanovic, M. Tallmadge, DH. Eberhart, BL. Ewald, DK. Gruber, DC. Morsch, TR. Strothers, MA, Rizzi, GP. Villigran, MD. (2003) Acrylamide Formation Mechanism in Heated Foods. *Journal of Agricultural and Food Chemistry* **51**, 4782-4787
- ¹⁵ Amrein TM., Bachmann S., Noti A., Biedermann M., Barbosa MF., Biedermann-Brem S., Grob K., Keiser A., Realini P., Escher F., Amado R. (2003) Potential of acrylamide formation, sugars and free asparagine in potatoes: A comparison of cultivars and farming systems. *Journal of Agricultural and Food Chemistry* **51**, 5556-5560
- ¹⁶ Stadler, R.H. Verzeegnassi, L. Varga, N. Grigorov, M. Studer, A. Riediker, S. Schilter, B. (2003) Formation of vinylogous compounds in model Maillard reaction systems. *Chemical Research and Toxicology* **16**, 1242-1250

- ¹⁷ Yasuhara, A; Tanaka, Y; Hengel, M and Shibamoto, T (2003) Gas Chromatographic Investigation of Acrylamide Formation in Browning Model Systems. *Journal of Agricultural and Food Chemistry* **51**, 3999-4003
- ¹⁸ Friedman, M., (2003) Chemistry, biochemistry, and safety of acrylamide. A review. *Journal of Agricultural and Food Chemistry*, **51**, 4504-4526.
- ¹⁹ Vatter, D. & Shetty, K (2003) Acrylamide in Food: a model for mechanism of formation and its reduction. *Innovative Food Science and Emerging Technologies* **4** (3), 331-338
- ²⁰ Ahn, JS. & Castle, L. (2003) Tests for the depolymerization of polyacrylamides as a potential source of acrylamide in heated foods. *Journal of Agricultural and Food Chemistry*. **51**, 6715-6718
- ²¹ Data not yet published.
- ²² Biedermann, M., Biedermann-Brem, S., Noti, A. and Grob K. (2002) Methods for determining the potential of acrylamide formation and its elimination in raw materials for food preparation, such as potatoes. *Mitteilungen aus Lebensmitteluntersuchung und Hygiene* **93**, 653-667.
- ²³ Rydberg, P. Eriksson, S. Tareke, E. Karlsson, P. Ehrenberg, L. Tornqvist, M. (2003) Investigations of factors that influence the acrylamide content of heated foodstuffs. *Journal of Agricultural and Food Chemistry* **51** (24) 7012-7018
- ²⁴ Taeymans, D.; Ashby, P.; Blank I., Gonde P.; Van Eijck, P.; Lalljie, S.; Lingnert, H.; Lindblom, M.; Matissek R.; Muller D.; O'Brien J.; *et al* (2003) A Review of Acrylamide: An Industry Perspective on Research, Analysis, Formation & Control. *Critical Reviews in Food Science & Nutrition*, **Submitted**.
- ²⁵ Ahn JS., Castle L, Clarke A., Lloyd M., Speck D. (2002) Verification of the findings of acrylamide in heated foods. *Food Additives and Contaminants* **19**, 1116-1124
- ²⁶ Grob, K., Biedermann, M., Biedermann-Brem, S., Noti, A., Imhof, D., Amrein, Th., Pfefferle, A., Bazzocco, D. (2003) French fries with less than 100 µg/kg acrylamide. A collaboration between cooks and analysts. *Eur. Food Res. Technol.* **217**, 185-194.
- ²⁷ Biedermann-Brem, S., Noti, A., Grob, K., Imhof, D., Bazzocco, D. and Pfefferle A. (2003) How much reducing sugar may potatoes contain to avoid excessive acrylamide formation during roasting and baking? *Eur. Food Res. Technol.* **217**, 369-373.
- ²⁸ Yung, M.Y.; Choi, D.S.; Ju, J.W. (2003) A novel technique for limitation of acrylamide formation in fried and baked corn chips and in French fries. *Journal of Food Science* **68** (4), 1287-1290
- ²⁹ European Parliament and Council Directive on Food additives other than Colours and Sweeteners (95/2/EC) (1995): http://europa.eu.int/comm/food/fs/sfp/addit_flavor/additives/index_en.html. Annex III, Part B.
- ³⁰ Springer, M.; Tischer, T.; Lehrack, A. & Freund, W. (2003) Development of acrylamide in bakery products. *Getreide Mehl und Brot* **57** (5), 274-278
- ³¹ Putz, B. (1993) The sugars of the potato as a subject of tuber-physiology. *European Association for Potato research: Abstracts of the 12th Triennial Conference, Paris, 1993*, 244-246.
- ³² Noti, A., Biedermann-Brem, S., Biedermann, M., Grob, K., Albisser, P., and Realini, P. (2003) Storage of potatoes at low temperature should avoided to prevent increased acrylamide formation during frying or roasting. *Mitteilungen aus Lebensmitteluntersuchung und Hygiene* **94**, 167-180.
- ³³ EC's Institute for Reference Materials and Measurements (2003) Acrylamide Monitoring Database: <http://www.irmm.jrc.be/ffu/acrylamide.html>

- ³⁴ Acrylamide Information Base of Research Activities in the EU:
http://europa.eu.int/comm/food/food/chemicalsafety/contaminants/acryl_database_en.htm
- ³⁵ FAO/WHO Acrylamide Infonet: <http://www.acrylamide-food.org/>
- ³⁶ EC 6th Framework Programme: http://europa.eu.int/comm/research/fp6/index_en.html